



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [info.bmjopen@bmj.com](mailto:info.bmjopen@bmj.com)

# BMJ Open

## Exploring the potential for introducing home monitoring of blood pressure during pregnancy into maternity care: current views and experiences of staff. A qualitative study.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2020-037874
Article Type:	Original research
Date Submitted by the Author:	19-Feb-2020
Complete List of Authors:	Hinton, Lisa; University of Cambridge, THIS Institute, Department of Public Health and Primary Care Hodgkinson, James ; University of Birmingham, Primary Care Clinical Sciences, Institute of Applied Health Research Tucker, Katherine; University of Oxford, Nuffield Department of Primary Care Rozmovits, Linda Chappell, Dr Lucy; King's College London, Women's Health Academic Centre Greenfield, Sheila; University of Birmingham, Primary Care Clinical Sciences McCourt, Christine; City University of London, Department of Midwifery and Child Health Sandall, Jane; Kings College, London, Health and Social Care Research Division McManus, Richard; University of Oxford, Dept of Primary Care Health Sciences
Keywords:	Hypertension < CARDIOLOGY, Organisation of health services < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Maternal medicine < OBSTETRICS, QUALITATIVE RESEARCH

SCHOLARONE™  
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

**Exploring the potential for introducing home monitoring of blood pressure during pregnancy into maternity care: current views and experiences of staff. A qualitative study.**

Hinton L<sup>1</sup>, Hodgkinson J<sup>2</sup>, Tucker K<sup>1</sup>, Rozmovits L<sup>4</sup>, Chappell LC<sup>3</sup>, Greenfield S<sup>2</sup>, McCourt C<sup>5</sup>, Sandall J<sup>3</sup>, McManus RJ<sup>1</sup>.

<sup>1</sup> Nuffield Department of Primary Care Health Sciences, University of Oxford OX2 6GG, UK.

<sup>2</sup> Institute of Applied Health Research, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK.

<sup>3</sup> Department of Women and Children's Health Academic Centre, King's College London, London, SE1 7EH

<sup>4</sup> Montreal, Quebec H2L 3X1, Canada

<sup>5</sup> Centre for Maternal and Child Health, School of Health Sciences, City University, London EC1R 1UW, UK

**Running title:** Staff views on home BP monitoring during pregnancy

**Author responsible for correspondence:** Lisa Hinton

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Tel +44 (0)1865 289300    lisa.hinton@phc.ox.ac.uk

Department of Primary Health Care Sciences, New Radcliffe House, Woodstock Rd, Oxford,  
  
UK, OX2 6GG

Word Count: 3,541

Keywords: hypertension, pregnancy, self-monitoring blood pressure, qualitative, maternity  
  
staff

## ABSTRACT

**Objective** One in 20 women are affected by pre-eclampsia, a major cause of maternal and perinatal morbidity, death and premature birth worldwide. Diagnosis is made from monitoring blood pressure (BP) and urine and symptoms at antenatal visits after 20 weeks of pregnancy. There are no randomized data from contemporary trials to guide the efficacy of self-monitoring of blood pressure (SMBP) in pregnancy. We explored the perspectives of maternity staff to understand the context and health system challenges to introducing and implementing SMBP in maternity care, ahead of undertaking a trial.

**Design.** Exploratory study

**Setting.** Eight hospitals, English National Health Service.

**Participants** Obstetricians, community and hospital midwives, pharmacists, trainee doctors (n=147)

**Methods** Semi-structured interviews with site research team members and clinicians, interviews and focus group discussions.

**Results** The main themes to emerge around SMBP include i) different blood pressure changes in pregnancy ii) reliability and accuracy of BP monitoring iii) anticipated impact of SMBP on women iv) anticipated impact of SMBP on the antenatal care system, v) caution, uncertainty and evidence vi) concerns over action/inaction and patient safety.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Conclusions** The potential impact of SMBP on maternity services is profound although nuanced. While introducing SMBP does not reduce the responsibility clinicians have for women’s health, it may enhance the responsibilities and agency of pregnant women, and introduces a new set of relationships into maternity care. This is a new space for reconfiguration of roles, mutual expectations, and the relationships between and responsibilities of health care providers and women.

## Article Summary

### Strengths and limitations of this study

- Qualitative study to explore maternity staff perspectives on self-monitoring of blood pressure in pregnancy
- Diverse sample includes voices from across multidisciplinary maternity teams
- Focus groups held across 8 hospitals in the English National Health Services, including hospital and community midwives



1  
2  
3  
4 **MAIN BODY OF TEXT**

5  
6  
7 **Introduction**

8  
9  
10 Raised blood pressure (BP) in pregnancy affects around one in ten women, almost half of  
11  
12  
13 whom develop pre-eclampsia, a major cause of maternal and perinatal morbidity, death and  
14  
15  
16 premature birth worldwide.(1, 2) Diagnosis is made from monitoring BP and urine and  
17  
18  
19 symptoms at antenatal visits after 20 weeks of pregnancy.  
20  
21  
22

23  
24  
25  
26 Self-monitoring of BP (SMBP) in the general population has become more common and large  
27  
28  
29 trials have demonstrated its effectiveness in terms of BP control. (3) Since 2011 UK national  
30  
31  
32 guidance for the general population with hypertension there has been a growing acceptance  
33  
34  
35 by clinicians of using SMBP values. Primary care physicians report an increasing use of SMBP  
36  
37  
38 to diagnose hypertension and for the ongoing management of hypertension to monitor BP  
39  
40  
41 control. (4-7)  
42  
43  
44  
45  
46  
47

48  
49 But what of the potential for introducing SMBP for women during or after pregnancy, to  
50  
51  
52 improve the detection and management of gestational hypertension and pre-eclampsia?  
53  
54  
55 Pregnant women are different from the general population: BP can rise rapidly during  
56  
57  
58 pregnancy, and the problems associated with this (such as pre-eclampsia, eclampsia and  
59  
60

stroke) are potentially serious. Pilot work in the UK suggests, with support from midwives and doctors, it is feasible and acceptable for women to monitor their BP and urine safely, potentially identifying hypertension earlier and controlling BP better.(8-10) SMBP during pregnancy is becoming common in some settings. A Canadian pilot study found more than 60% of women diagnosed with non-proteinuric hypertension in pregnancy were already undertaking SMBP. (11) The Society of Obstetricians and Gynaecologists of Canada provides guidelines for SMBP in pregnancy and most Canadian obstetricians and primary care physicians used SMBP to assess for white coat hypertension (WCH), (12) but its place in other countries is not yet clear. (13)

While SMBP in pregnancy is already implemented in some UK settings, (14) there are no adequately powered randomized data from contemporary trials to guide the efficacy of such methods. Our early work exploring women's experiences of taking part in an SMBP pilot study (15) showed that most found their health professionals to be supportive, although in some cases self-monitoring results were ignored. An understanding of the attitudes of individuals and the organization will be paramount in introducing new, and safe, models of care. Ahead of undertaking a large trial of BP monitoring in high-risk pregnancy to improve detection and monitoring of hypertension, we undertook a study to explore the perspectives of staff across

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

the maternity care pathway to explore the context and health system challenges to introducing and implementing SMBP in maternity care. (10, 16, 17)

Methods

We carried out an intervention development phase in preparation for a large trial of SMBP in pregnancy, aiming to understand the views and prior experience of clinicians regarding such monitoring. This involved interviews and focus groups conducted in 2016-2017 across eight participating teaching and urban general district hospitals that serve diverse populations. (TABLE 1). We conducted face-to-face interviews with a lead consultant obstetrician at each site (n=8) and a series of focus groups and interviews with a wide range of healthcare professionals.

**Sample/setting** We sought a broad pragmatic sample that included community and hospital-based midwives and obstetricians of different seniorities at each site (n=147). All focus groups and interviews took place at hospital or health centre sites, either as part of, or after, existing meetings or during shift breaks. It was usually a Research Midwife at each site who identified who might be available with some of the larger focus groups piggy-backing on existing meetings. Focus group size ranged from two to 15 individuals.

**Data collection** The interviews and focus groups were facilitated by two social science/non-clinical researchers (LH and JH, male and female) and audio-recorded for transcription and

analysis. A topic guide was developed drawing on the literature, our prior work and in discussion with the multi-disciplinary trial team. Questions sought to explore issues around staff views on SMBP in pregnancy, the accuracy of home readings, how to manage care of women who present with home readings (particularly those discordant with clinic readings), the impact of home readings on clinical decision-making and on workload, as well as practical issues around operationalising and integrating self-monitoring into current care pathways.

**Analysis** All interviews and discussions were transcribed for analysis, which was conducted in two phases. An initial rapid analysis was undertaken to guide trial development. (18) This used templates developed by JH and LH based on the topic guides and subsequently refined after a period of piloting. We conducted a rapid content analysis to identify evidence related to the a priori concepts and emergent themes that would then guide development of the study. In a subsequent phase LH and JH, in discussion with the wider team, developed a coding frame and worked in parallel (with LR) to code and analyse the data.(19) Thematic analysis was undertaken to explore the anticipated and emergent themes and was then mapped back to the framework, supported by NVIVO (20, 21). Data collection continued until data saturation was reached for the themes reported in this paper. (22) Illustrative quotes are presented in TABLE 2. We used SPQR checklist when writing our report(23).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Patient and public involvement** We have patient and public involvement for the overall programme of research, but as this was a staff focused study, patients were not directly involved in the phase of work reported here.

RESULTS

Our findings suggest SMBP during pregnancy was not yet a widespread practice across the sites included. However, some midwives and obstetricians reported occasional pregnant women who were already self-monitoring, either of their own volition or because they had been advised to because of previous experiences of pre-eclampsia, white coat hypertension (WCH), or pre-existing hypertension. The main themes to emerge around SMBP were i) different blood pressure changes in pregnancy ii) reliability and accuracy of BP monitoring iii) anticipated impact of SMBP on women iv) anticipated impact of SMBP on the antenatal care system, v) caution, uncertainty and evidence vi) concerns over action/inaction and patient safety.

*Different blood pressure changes in pregnancy*

BP fluctuation

Staff emphasised that BP is different in pregnancy than at other times in women’s lives. It fluctuates throughout pregnancy and these fluctuations are normal. However, BP in

pregnancy can change rapidly, precipitating a medical emergency for mother and baby.

While antenatal care staff will be familiar with this phenomenon through their training and experience, women may not be. There was concern women could have difficulty interpreting results and deciding when to seek acute care.

In the general population, BP readings are used to guide care and medication changes over the long term, but in pregnancy the situation is more dynamic. Some were interested in the wider context home BP readings would give them, in particular, feeling women with WCH might get more typical readings at home. However, several indicated they would tend to privilege their own clinic readings as they reflected the situation happening immediately in front of them.

### Wider symptoms

Staff were also concerned that BP is not the only factor in diagnosing hypertension or pre-eclampsia but rather one piece of a diagnostic 'jigsaw'. Healthcare professionals stressed the importance of ensuring women were alert to the wide range of symptoms (e.g. headaches, dizziness, blurred vision, proteinuria) and would act on those even if their BP appeared to be in the normal range.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*Reliability and accuracy*

The reliability and accuracy of machines were key concerns for staff, as well as whether women would monitor their BP correctly and respond appropriately. Many raised concerns about the quality of home BP monitors, whether women were using one of the relatively few validated for use in pregnancy, and whether they were properly calibrated. There were concerns about appropriate cuff size, particularly for women with a high body mass index. Other factors identified as potentially impacting the reliability and accuracy of home readings included women’s practices and the context in which they took readings (did they sit correctly, choose a quiet time of day?).

*Impact on women*

Staff had a range of perceptions about how pregnant women might feel about being asked to self-monitor their BP. Some anticipated women being happy to participate and enjoying a sense of agency, control and reassurance over their own healthcare. Others raised concern that women may experience increased anxiety and become obsessed about their BP. Some felt pregnant women expect care to be led by providers and asking them to monitor their own BP would be perceived as a way of abdicating responsibility. Although staff also

acknowledged the potential for empowerment, some expressed concern that asking women to self-monitor would contribute to an over-medicalisation of pregnancy in contrast with the evolution of antenatal care away from this model in recent years.

Many potentially positive impacts for women were identified both in terms of quality of experience and better outcomes for mother and baby. These included fewer medical appointments, shorter waiting times and avoiding hospitalisation. Staff acknowledged detecting BP increases earlier would have the potential to deliver better outcomes for both mothers and babies, but confirmed definitive evidence of these outcomes would be needed before they could commit to changing their practice.

### *Anticipated impact on antenatal care*

Staff were caught between feeling SMBP could reassure women, so cutting 'worried well' visits and/or raise anxieties, thus increasing them. So, while they recognised other potential health system benefits including fewer women coming into clinic, thus freeing up clinic time in hospital and in community care, they also raised various concerns around the impact of SMBP on antenatal care pathways and their workload. These included a rise in phone calls to maternity assessment units and an increased demand for care, particularly from the 'worried well'.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Staff were aware that, ahead of definitive trial results, these reflections were hypothetical.

Their currently available best comparator was their experience with women’s self-monitoring for gestational diabetes, which is now quite common. Some described SMBP as a challenge to their professional roles, particularly for midwives, and part of wider changes towards increasing self-care and personal responsibility in health. One consultant described the shifting responsibilities in antenatal care, with health education now as much a part of routine care as systematic screening for obstetric complications.

*Caution, uncertainty and evidence*

While staff were generally open to looking at home BP readings, most were mindful the evidence was not yet available as to how to treat these readings in conjunction with clinic readings. Most felt they could only treat home readings as additional rather than as core information, not directing care decisions, but contributing extra detail to the ‘jigsaw’:

**Caution**

Many midwives said that if self-monitoring readings were discrepant with clinic readings, they would privilege the readings they had just taken themselves. Most added they would take into

1  
2  
3 account the home readings, though exactly how would be considered on a case-by-case  
4  
5  
6 basis. Some would use them as part of improving understanding of the wider context, but use  
7  
8  
9 their clinic reading for the decision on whether to admit a woman to hospital. The home reading  
10  
11  
12 would not be used for treatment decisions but could influence how often an obstetrician felt it  
13  
14  
15 necessary to see a woman.  
16  
17

18  
19  
20  
21  
22 Some acknowledged the potential of SMBP to identify WCH. But in several midwife  
23  
24  
25 discussions, consensus emerged on the side of caution, such as referring women onwards or  
26  
27  
28 taking bloods if only one type of reading (clinic or home) was high. Others referred to local  
29  
30  
31 protocols and guidelines although this was often interpreted as needing to act on any higher  
32  
33  
34 reading, implying caution and a focus on patient safety, with fear of being considered negligent  
35  
36  
37 explicitly raised by one midwife.  
38  
39  
40

41  
42  
43  
44  
45 Some said they verified home readings by testing the device used by women alongside their  
46  
47  
48 clinic monitor. Several midwives commented they would have to respect a woman's readings,  
49  
50  
51 as to do otherwise could affect mutual trust.  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Uncertainty

There was awareness that pre-eclampsia is a variable condition in which the speed of disease progression varies considerably. Some felt SMBP could significantly mitigate some of these uncertainties. One senior obstetrician specifically alluded to a potential *advantage* of SMBP being its ability to reduce uncertainty by enabling clinicians to make objective assessment of how accurate clinic readings are, clarifying who has WCH, assessing the effectiveness of treatment and standardising the various clinical environments that women might be seen in. Yet others clearly feared SMBP could increase uncertainty (discrepancy in readings), and/or lead to overcaution and overtreatment on the part of less experienced colleagues.

Different approaches to managing hypertension in pregnancy also emerged. While some make judgements on a case-by-case basis, privileging the extent of change from baseline, the wider context, and individualised pathways, and given that different hospitals may have their own guidelines, others described a practice defined by the National Institute for Health and Care Excellence (NICE) guidelines and the use of thresholds as the defining guide in when to intervene (24).

## Evidence

Numerous staff raised the need for more and improved evidence. One obstetrician observed that guidelines are supposed to be evidence based but recommendations are often made on expert opinion. A different senior obstetrician also suggested that people practise defensively, leading to many unnecessary blood tests being done because a greater number of healthcare professionals (midwives and junior doctors) are able to request the tests, and because when women have been told they may have pre-eclampsia, heightened anxiety leads to repeated checks for reassurance. Overall, there was a distinct lack of consensus, but with a clear wish from many for more evidence to reduce uncertainty.

## *Concerns over action/inaction*

While a shift to SMBP is in keeping with wider trends towards self-care and personal responsibility, working through this new balance of responsibility for acting on readings was a cause for concern. Staff raised concerns about women not acting on readings, through either not realising they required urgent action, not wanting to bother busy staff, or not wanting to be hospitalised. There were concerns about women ignoring problems, or being falsely reassured by one reading and ignoring the wider context. Many stressed the importance of women

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

having a clear understanding of what they were supposed to do if their readings were high, with a clear pathway for women presenting with high BP based on home readings.

Many staff made comparisons with home monitoring of blood glucose in gestational diabetes, either as a useful precedent for SMBP or evidence of the challenges of women deliberately or inadvertently coming in with inaccurate results.

Others expressed the view that it is hospital policy (as well as being the health professional's view) that they would rather ask women to come in for comprehensive assessment than have the woman check her BP at home and decide not to come in.

DISCUSSION

Main findings.

Antenatal care staff were generally in favour of asking women to monitor their own BP at home, but flagged concerns about the potential impacts on women and staff workload, the reliability and accuracy of readings and the need for clarity about how the information would be used by clinicians. Many emphasised the importance of ensuring women understood what to do in the event of readings that cause concern and provided examples of other situations where women had failed to act in acute situations because they did not understand the severity

1  
2  
3 of the situation or rationalised away the need to seek medical attention. Some voiced concerns  
4  
5  
6 about women becoming either anxious or falsely reassured by self-monitoring. There was a  
7  
8  
9 feeling that SMBP might only be for a select few as it will require a shift in thinking about who  
10  
11  
12 takes responsibility for the woman's health.  
13  
14  
15  
16  
17  
18

19 Many of the issues raised were specific to pregnancy. However, concerns about the potential  
20  
21  
22 for monitoring leading to anxiety may also exist for the general population – though previous  
23  
24  
25 studies have not found evidence to support this (25) – they could be intensified when there  
26  
27  
28 may be additional anxiety related to the effect of BP changes on the baby. Concerns about  
29  
30  
31 device accuracy were well founded as relatively few monitors on the market are validated in  
32  
33  
34 pregnancy.(26) The need to consider more than just BP and the dynamic nature of BP in  
35  
36  
37 pregnancy as opposed to BP in the general population means that failure to act has the  
38  
39  
40 potential for devastating consequences for both woman and baby. Yet resistance to  
41  
42  
43 medicalisation may be heightened in a previously healthy younger population of 'women' as  
44  
45  
46 opposed to 'patients'. (27) Continuity of care is an ambition not always achieved in maternity  
47  
48  
49 care. The lack of an established relationship of trust between women/patients and healthcare  
50  
51  
52 professionals that is typical of hypertension management in primary care, further complicates  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

the situation (28). Staff are therefore understandably cautious and uncertain, while also willing to see the potential for SMBP to reduce uncertainty and add usefully to available evidence.

Strengths and limitations

A strength of this study is the breadth of multidisciplinary voices across the maternity care pathway captured, spanning eight differently located hospitals in England. This has given rich insights into the potential challenges of introducing SMBP into maternity care from the perspectives of different disciplines and locales. Our two-stage approach to analysis, led by social scientists with expert clinical input from co-authors in obstetrics, midwifery and primary care, has also enhanced the interpretive strength of the findings. Limitations include the challenges of holding focus group discussions with busy professionals engaged with frontline care. These are vital perspectives to capture but there were inevitably challenges in terms of finding time and space for some focus group discussions.

There are several literatures to draw on in interpreting these results.

**Responsibilisation:**To date, measuring and monitoring women’s BP in pregnancy has been a central component of the healthcare professional’s role in antenatal care. To include women’s home readings into these care pathways represents a significant shift in the balance of responsibility. Emerging literature that explores the shifts and implications of health

1  
2  
3 'responsibilisation' are of relevance. Shifts in the balance of responsibility have the potential  
4  
5  
6  
7 to disrupt long established care pathways and can be experienced both positively as well as  
8  
9  
10 negatively as surveillance.(29) Exploring the impacts of these shifts is a live issue in the  
11  
12  
13 sociological literature from studies of 'park runs' to HIV management (30, 31), and the  
14  
15  
16 relationships between social workers and vulnerable youth .(32) Drawing on work by Garret  
17  
18  
19 (2008), Liebenberg et al explored the impact of shifts in neo-liberal systems, where individuals  
20  
21  
22 are expected to manage their own risks and demonstrate self-care. "*These shifts impact on*  
23  
24  
25 *the roles that health professionals play, from active management to encouraging/teaching of*  
26  
27  
28 *the skills of management*" (33). Newman et al's 2012 study of young people with HIV revealed  
29  
30  
31 narratives of responsibilisation that can give rise to contest between young people and their  
32  
33  
34 clinicians. (31) In exploring the impact of telecare on the management of long term conditions,  
35  
36  
37 Rogers et al (2011) concluded "*Indeed a paradox of the reliance and acceptance of telecare*  
38  
39  
40 *is the creation of new relationships and dependencies rather than the diminution of reliance*  
41  
42  
43 *envisaged by policy.*" (34) Our findings on clinician caution and concerns over inaction,  
44  
45  
46 indicate that these shifts in responsibility could have profound impacts that we need to be alive  
47  
48  
49  
50  
51 to as we further explore the impact of this work going forward. (35)  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Impact on professional roles:** A shift in responsibility for BP monitoring may also have an impact on professional roles and identity. Concern over these impacts ran through various themes we have reported, including impact on workload, caution and uncertainty and location of expertise. Task shifting is a key issue in many healthcare settings, but it does not emerge seamlessly as revealed in the literature.(36-39) How SMBP impacts on these professional roles, and how professionals behave when there is a shift in the locus of responsibility will need to be the focus for future work.

**Patient safety:** While staff raised concerns about the impact of SMBP on patient safety, these concerns were expressed in the context of an emerging evidence base for SMBP (13, 17). As this develops, we draw on evidence that points to the considerable scope for improving patient and family contributions to the detection and management of acute illness (40). Delayed recognition and treatment of conditions such as pneumonia and meningitis in childhood (41), pre-eclampsia and reduced fetal movements during pregnancy and after childbirth (42, 43) and heart disease and stroke in adulthood, (44, 45) contribute significantly to the mortality and morbidity burden. These conditions typically present with a time-critical window for early recognition and response, and are associated with red flag signs and symptoms (such as breathlessness and pain) which can signify a serious underlying condition and act as potential

1  
2  
3 markers to aid patient and family involvement in escalation of care. However, research has  
4  
5  
6 shown there are challenges to speaking up and raising safety alerts in maternity care and that  
7  
8  
9 organization-focused efforts are required to improve staff responsiveness. (46)  
10  
11  
12  
13  
14  
15

## 16 Conclusions

17 We need to be mindful that the impact of SMBP on maternity services could be profound  
18  
19 although nuanced. Whilst its introduction will not reduce the responsibility clinicians have for  
20  
21  
22 women's health, it promises to enhance the responsibilities and agency of women, and  
23  
24  
25 introduces a new set of relationships into maternity care. We cannot simply look to the  
26  
27  
28 literature and experiences of SMBP in other populations (e.g. such as hypertension outside  
29  
30  
31 of pregnancy) because of the very nature of BP changes in pregnancy and because this is a  
32  
33  
34 healthy population, unless specifically indicated otherwise. This is, then, a new space for  
35  
36  
37 reconfiguration of roles, mutual expectations, and the relationships between, and  
38  
39  
40 responsibilities of, health care providers and mothers.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

## 53 ACKNOWLEDGEMENTS

54  
55  
56 We would like to thank all NHS staff who gave their time to contribute to this study.  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

DISCLOSURE OF INTERESTS

RJM has received BP monitors for research from Omron. No other authors have conflicts to declare.

DATA STATEMENT

Transcripts are held by the University of Oxford.

AUTHOR CONTRIBUTIONS

Authors contributions are as follows: LH, JH, KT, LCC, SG, CM, JS, LR and RM conceived the study, secured the funding and were involved in the planning and carrying out the study. LH and JH conducted the interviews and focus groups and KT managed the study. LH, JH and LR analysed the data with input from the wider authorship. LH and JH drafted the paper. All authors commented on drafts and writing up the work.

DETAILS OF ETHICS APPROVAL

A favourable ethical review for this study was obtained from South Central - Oxford C Research Ethics Committee (16/SC/0386). Written informed consent was gained from individuals who agreed to take part.

## FUNDING

This article represents independent research commissioned by the National Institute for Health Research (NIHR) Programme for Applied Research (RP-PG-0614-20005). Richard McManus was and Lucy Chappell is supported by a Research Professorship from the National Institute for Health Research (NIHR-RP-R2-12-015 and RP-2014-05-019 respectively). Richard McManus is an NIHR Senior Investigator. Lisa Hinton, Katherine Tucker and Richard McManus have received funding from the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care Oxford at Oxford Health NHS Foundation Trust and NIHR Oxford Thames Valley Applied Research Collaboration. Lisa Hinton is also supported by the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC). Jane Sandall is a National Institute for Health Research (NIHR) Senior Investigator and is supported by the National Institute for Health Research (NIHR) Applied Research Collaboration South London (NIHR ARC South London) at King's College Hospital NHS Foundation Trust. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health.

REFERENCES

1. Mol BW, Roberts CT, Thangaratinam S, Magee LA, De Groot CJ, Hofmeyr GJ. Pre-eclampsia. *The Lancet*. 2016;387(10022):999-1011.

2. Conti-Ramsden F, Knight M, Green M, Shennan AH, Chappell LC. Reducing maternal deaths from hypertensive disorders: learning from confidential inquiries. *Bmj*. 2019;364:l230.

3. McManus R, Wood S, Bray E, Glasziou P, Hayen A, Heneghan C, et al. Self-monitoring in hypertension: a web-based survey of primary care physicians. *Journal of human hypertension*. 2014;28(2):123.

4. National Clinical Guideline Centre. Hypertension: management of hypertension in adults in primary care. London: National Institute for Clinical Excellence; 2011.

5. McManus RJ, Mant J, Bray EP, Holder R, Jones MI, Greenfield S, et al. Telemonitoring and self-management in the control of hypertension (TASMINH2): a randomised controlled trial. *The Lancet*. 2010;376(9736):163-72.

6. McManus RJ, Mant J, Franssen M, Nickless A, Schwartz C, Hodgkinson J, et al. Efficacy of self-monitored blood pressure, with or without telemonitoring, for titration of antihypertensive medication (TASMINH4): an unmasked randomised controlled trial. *The Lancet*. 2018;391(10124):949-59.

7. Fletcher BR, Hinton L, Bray EP, Hayen A, Hobbs FR, Mant J, et al. Self-monitoring blood pressure in patients with hypertension: an internet-based survey of UK GPs. *Br J Gen Pract*. 2016;66(652):e831-e7.

8. Tucker KL, Taylor KS, Crawford C, Hodgkinson JA, Bankhead C, Carver T, et al. Blood pressure self-monitoring in pregnancy: examining feasibility in a prospective cohort study. *BMC pregnancy and childbirth*. 2017;17(1):442.

9. Tucker KL, Bowen L, Crawford C, Mallon P, Hinton L, Lee M-M, et al. The feasibility and acceptability of self-testing for proteinuria during pregnancy: A mixed methods approach. *Pregnancy hypertension*. 2018;12:161-8.

10. Mackintosh N, Rance S, Carter W, Sandall J. Working for patient safety: a qualitative study of women's help-seeking during acute perinatal events. *BMC pregnancy and childbirth*. 2017;17(1):232.

11. Magee LA, von Dadelszen P, Chan S, Gafni A, Gruslin A, Helewa M, et al. Women's views of their experiences in the CHIPS (Control of Hypertension in Pregnancy Study) Pilot Trial. *Hypertension in Pregnancy*. 2007;26(4):371-87.

12. Dehaeck U, Thurston J, Gibson P, Stephanson K, Ross S. Blood pressure measurement for hypertension in pregnancy. *Journal of Obstetrics & Gynaecology Canada: JOGC*. 2010;32(4):328-34.

13. Magee LA, Pels A, Helewa M, Rey E, von Dadelszen P. Diagnosis, evaluation, and management of the hypertensive disorders of pregnancy: executive summary. *J Obstet Gynaecol Can*. 2014;36(7):575-6.

14. Xydopoulos G, Perry H, Sheehan E, Thilaganathan B, Fordham R, Khalil A. Home blood-pressure monitoring in a hypertensive pregnant population: cost-minimization study. *Ultrasound in Obstetrics & Gynecology*. 2019;53(4):496-502.

15. Hinton L, Tucker KL, Greenfield SM, Hodgkinson JA, Mackillop L, McCourt C, et al. Blood pressure self-monitoring in pregnancy (BuMP) feasibility study; a qualitative analysis of women's experiences of self-monitoring. *BMC pregnancy and childbirth*. 2017;17(1):427.

16. Carter W, Bick D, Mackintosh N, Sandall J. A narrative synthesis of factors that affect women speaking up about early warning signs and symptoms of pre-eclampsia and responses of healthcare staff. *BMC pregnancy and childbirth*. 2017;17(1):63.

17. Dougall D. FM, Tucker K., Yu L-M., Hinton L., Rivero-Arias O., Abel L., Allen J., Band R., Chisholm A., Crawford C., Green M., Greenfield S, Hodgkinson J., Leeson P., McCourt C., Mackillop L., Nickless A., Sandall J., Santos M., Tarassenko L., Velardo C., Wilson H., Yardley L., Chappell LC., McManus RJ. . Blood pressure monitoring in high risk pregnancy to improve the detection and monitoring of hypertension (The BUMP 1&2 trials): protocol for two linked randomised controlled trials. *BMJ Open*. 2019.
18. Hamilton A. Qualitative Methods in Rapid TurnAround Health Services Research VA HSR&D Cyberseminar Spotlight on Women's Health 2013 26.06.2017. Available from: [https://www.hsr.d.research.va.gov/for\\_researchers/cyber\\_seminars/archives/780-notes.pdf](https://www.hsr.d.research.va.gov/for_researchers/cyber_seminars/archives/780-notes.pdf).
19. Gale NK, Heath G, Cameron E, Rashid S, Redwood S. Using the framework method for the analysis of qualitative data in multi-disciplinary health research. *BMC medical research methodology*. 2013;13(1):117.
20. Ziebland S, McPherson, A. Making sense of qualitative data analysis: an introduction with illustrations from the DIPEX (personal experiences of health and illness. *Medical Education*. 2006;40(5):405-14.
21. QSR. NVIVO 9 Qualitative data analysis software. 2018.
22. Mason M, editor Sample size and saturation in PhD studies using qualitative interviews. *Forum Qualitative Sozialforschung/Forum: Qualitative Social Research*; 2010.
23. O'Brien BC, Harris IB, Beckman TJ, Reed DA, Cook DA. Standards for reporting qualitative research: a synthesis of recommendations. *Acad Med*. 2014;89(9):1245-51.
24. National Institute for Health and Care Excellence N. Hypertension in pregnancy: diagnosis and management NICE guideline [NG133]. 2019.
25. Jones MI, Greenfield SM, Bray EP, Baral-Grant S, Hobbs FR, Holder R, et al. Patients' experiences of self-monitoring blood pressure and self-titration of medication: the TASMINH2 trial qualitative study. *British Journal of General Practice*. 2012;62(595):e135-e42.
26. Hodgkinson JA, Tucker KL, Crawford C, Greenfield SM, Heneghan C, Hinton L, et al. Is self monitoring of blood pressure in pregnancy safe and effective? *BMJ*. 2014;349:g6616.
27. Johanson R, Newburn M, Macfarlane A. Has the medicalisation of childbirth gone too far? *BMJ*. 2002;324(7342):892-5.
28. McCourt C, Stevens T. Continuity of carer—what does it mean and does it matter to midwives and birthing women? *Canadian Journal of Midwifery Research and Practice*. 2006;4(3):10-20.
29. Bunton R, Petersen, A. Foucault, Health and Medicine. London: Routledge; 2002.
30. Wiltshire G, Fullagar S, Stevenson C. Exploring parkrun as a social context for collective health practices: running with and against the moral imperatives of health responsibilisation. *Sociology of health & illness*. 2018;40(1):3-17.
31. Newman CE, Persson A, Miller A, Brown RJ. "Just take your medicine and everything will be fine": Responsibilisation narratives in accounts of transitioning young people with HIV into adult care services in Australia. *AIDS care*. 2016;28(1):131-6.
32. Liebenberg L, Ungar M, Ikeda J. Neo-liberalism and responsibilisation in the discourse of social service workers. *The British Journal of Social Work*. 2013;45(3):1006-21.
33. Garrett PM. How to be modern: New Labour's neoliberal modernity and the Change for Children programme. *British Journal of Social Work*. 2006;38(2):270-89.
34. Rogers A, Kirk S, Gately C, May CR, Finch T. Established users and the making of telecare work in long term condition management: implications for health policy. *Social science & medicine*. 2011;72(7):1077-84.
35. Cohn S. From health behaviours to health practices: an introduction. *Sociology of health & illness*. 2014;36(2):157-62.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

36. May C, Finch T, Mair F, Ballini L, Dowrick C, Eccles M, et al. Understanding the implementation of complex interventions in health care: the normalization process model. *BMC health services research*. 2007;7(1):148.

37. Ferrinho P, Sidat M, Goma F, Dussault G. Task-shifting: experiences and opinions of health workers in Mozambique and Zambia. *Human Resources for Health*. 2012;10(1):34.

38. Dawson AJ, Buchan J, Duffield C, Homer CS, Wijewardena K. Task shifting and sharing in maternal and reproductive health in low-income countries: a narrative synthesis of current evidence. *Health policy and planning*. 2013;29(3):396-408.

39. Bohmer RM, Imison C. Lessons from England's health care workforce redesign: no quick fixes. *Health Affairs*. 2013;32(11):2025-31.

40. Mackintosh NJ. Interventions to increase patient and family involvement in escalation of care for acute life-threatening illness in community health and hospital settings. 2017.

41. Wolfe I, Cass H, Thompson MJ, Craft A, Peile E, Wieggersma PA, et al. Improving child health services in the UK: insights from Europe and their implications for the NHS reforms. *Bmj*. 2011;342:d1277.

42. Draper ES, Kurinczuk JJ, Kenyon S. MBRRACE-UK Perinatal confidential enquiry: term, singleton, normally formed, antepartum stillbirth: Infant Mortality and Morbidity Studies; 2015.

43. Warland J, O'Brien LM, Heazell AE, Mitchell EA. An international internet survey of the experiences of 1,714 mothers with a late stillbirth: the STARS cohort study. *BMC pregnancy and childbirth*. 2015;15(1):172.

44. Schwappach DL. Engaging patients as vigilant partners in safety: a systematic review. *Medical Care Research and Review*. 2010;67(2):119-48.

45. Intercollegiate Stroke Working Party. Public Report for England, Wales and Northern Ireland. National Sentinel Stroke Clinical Audit 2010. 2010.

46. Rance S, McCourt C, Rayment J, Mackintosh N, Carter W, Watson K, et al. Women's safety alerts in maternity care: is speaking up enough? *NBMJ Quality & Safety*. 2013;22(4):348-55.



DATA TABLE 1.

HOSPITAL		NUMBERS	Staff
Guys & St Thomas's NHS Foundation Trust	PI Interview	1	Consultant Obstetrician
West Middlesex University Hospital	PI Interview	1	Consultant Obstetrician
The Royal London Hospital, Barts Health NHS Trust	PI Interview	1	Consultant Obstetrician
Whipps Cross Hospital, Barts Health NHS Trust	PI Interview	1	Consultant Obstetrician
The Royal Wolverhampton NHS Trust	PI Interview	1	Consultant Obstetrician
Birmingham Women's NHS Foundation Trust	PI Interview	1	Consultant Obstetrician
John Radcliffe Hospital, Oxford University Hospitals NHS Foundation Trust	PI Interview	1	Consultant Obstetrician
Royal Berkshire NHS Foundation Trust	PI Interview	1	Consultant Obstetrician
	<b>Total</b>	<b>8</b>	
HOSPITAL	FOCUS GROUP	NUMBERS	Staff represented
Guys & St Thomas's NHS Foundation Trust	Focus group 1	10	Obstetricians, midwives, pharmacist
	Focus group 2	4	Day assessment unit midwives
West Middlesex University Hospital	Focus group 1	15	Obstetricians, midwives, healthcare professionals
Whipps Cross Hospital, Barts Health NHS Trust	Focus group 1	11	Obstetricians, midwives
	Focus group 2	4	Antenatal midwives
	Focus group 3	4	Day assessment unit midwives
The Royal London Hospital, Barts Health NHS Trust	Focus group 1	6	Community midwives (FMU)
	Focus group 2	4	Community midwives (in hospital)
	Interview	1	Consultant obstetrician
	Focus group 3	7	Antenatal midwives
	Focus group 4	2	Medical trainees



John Radcliffe Hospital, Oxford University Hospitals NHS Foundation Trust	Focus group 1	11	Community midwives (in hospital)
	Focus group 2	3	Obstetricians
	Interview	1	Obstetrician
	Interview	1	Consultant obstetrician
	Interview	1	Obstetrician
	Focus Group 3	2	Hospital midwives (MAU)
	Interview	1	Hospital midwife (MAU)
Horton General Hospital, Oxford University Hospitals NHS Foundation Trust	Focus Group 4	13	Community midwives
Royal Berkshire NHS Foundation Trust	Focus group 1	2	Hospital midwives
	Focus group 2	2	Hospital midwives
	Interview	1	Hospital midwife
	Focus group 3	2	Obstetricians
	Interview	1	Obstetrician
	Focus group 4	6	Community midwives (in hospital)
The Royal Wolverhampton NHS Trust	Focus group 1	3	Obstetricians
	Focus group 2	2	Hospital midwives
	Focus group 3	10	Hospital and community midwives
	Focus group 4	8	Hospital and community midwives
	Interview	1	Hospital midwife
	<b>Total</b>	<b>139</b>	

TABLE 2 QUOTATIONS

THEME	QUOTATIONS
i) Different blood pressure changes in pregnancy	<p><b><i>BP Fluctuations</i></b></p> <p><i>"even as midwives we get confused with what should be normal because we have a protocol that they have a booking [.....measurement] so we've got a start and if it's 15 either way for the diastolic and we have to sort of think well yes that's creeping up a little bit if these girl's sent them to me I would look at it and say well that's not particularly high blood pressure because I'm dealing with PET all the time, you know, high blood pressure to me is 150/90 and above. So, isn't it, it's quite difficult and I think for a midwife let alone a woman to sit at home and think is that high am I going to be bothered"</i> <b>West Middlesex FG3</b></p> <p><i>"Another thing outside of pregnancy and not, most of what you'll doing with hypertension monitoring is trying to prevent long term complications on a short term and therefore, actually what you, slightly over 24 hours probably is what really matters whereas in pregnancy [...] I'm actually only interested in whether or not they've got [...] Well actually I'm really interested in the next 24 hours."</i> <b>Guys and St Thomas's FG1</b></p> <p><b><i>Wider symptoms</i></b></p> <p><i>"Yes but I think you raise a really important point about when you start to kind of focus on individual symptoms of a condition it's, there's a risk that you just focus on the blood pressure which means if you're feeling dizzy or your ankles are puffy or you're not really feeling yourself and got a blinding</i></p>

	<p><i>headache you don't focus on any those things because the blood pressure says it's okay."</i><b>Royal London, Community midwife</b></p> <p><i>"I think that they're over checking their blood pressure and occasionally I think it is helpful that they see a midwife and do things like check the protein in the urine and things like that which, you know, occasionally perhaps not be able to manage quite so well yourself"</i> <b>John Radcliffe, Obstetrician</b></p> <p><i>"So you kind of think oh are we only doing half of a job if they're just monitoring their blood pressure"</i> <b>Horton, Community Midwife</b></p>
ii) Reliability and accuracy	<p><i>"As a health professional I know blood pressure's affected by many things, whether you are taking it standing whether you are taking it lying whether it's in the morning whether you, or you've just come back from Tesco with plastic bags and stuff like, you're out of your breath or you've just fought with your partner or something like that, so many things influences blood pressure. So they will need to be given the empowerment they will need to take their blood pressure so that we, it's as accurate as, so that we get as close to the normal if that can be achieved, something like that."</i><b>West Middlesex FG2</b></p> <p><i>"[R4] . Because there used to be a particular, yes because women's bodies are totally different in pregnancy to somebody having a blood pressure normally and so, you know, for me I've never looked into it but whether it's a 100% accurate I don't know but I certainly don't like those box things because they have to be quiet, silent and not moving and nobody talking to them.</i></p> <p><i>[R1]: That's why the machine is important.</i></p>

[R4]: To get a really accurate result."Whipps Cross, FG1

*"I wouldn't ignore it if it was very elevated but I would certainly repeat it, I wouldn't go with what they said... Because I'd be worried about the quality of their machine"* John Radcliffe, Obstetrician

*"we advise them not to because [um] they're not calibrated, you know, they're their own there's no way they're calibrated and also quite a lot of [um] blood pressure [um] equipment, we've read research that says it's not as accurate as manual or a professional taking the blood pressure"* Royal Berkshire, FG1

*"when they come along and they bring readings [um] and they, if there's any conflict and sometimes they don't want to be labelled as hypertensive because it leads to certain choices later on in pregnancy that may be removed from them i.e. where they deliver, how they deliver [um] I then, you know, I will talk to them and say you, whatever you're using at home may or may not be validated for pregnancy but what we use here is validated for pregnancy and therefore I would defer to those readings"* Royal Wolverhampton, Consultant Obstetrician

*"I tend to be a little bit sceptical because you don't know what they're using, you don't know what machine they're using, you don't know whether they're trained to use it, you don't know how old that machine is whether it's again cuff size, you know, if they've borrowed it off somebody you don't know whether it's been PAT tested, you know, and if they are concerned about their reading being particularly high or particularly low I'd rather just get them in to see the relevant person at the time*

	<p><i>rather than rely on that, you know, I do worry about them using their own equipment because you just don't know how accurate it is"</i> <b>Royal Wolverhampton, FG2</b></p> <p><i>"I'd want to know whether or not the machine was up to scratch xxx good if she could correlate it with ours, bring it in we'll just make sure your machine is similar to ours, there's things like that."</i> <b>Royal Wolverhampton FG3</b></p>
iii) Impact on women	<p><i>"Those people with raised blood pressure will share their experience of like three or four weeks admission into hospital I think they will be delighted to do it to save coming in, they always, you know, get very upset having to stay on for three weeks and when they have child at home."</i> <b>West Middlesex FG3</b></p> <p><i>"So for the women who have had severe pre-eclampsia in the past who are absolutely panic stricken about it being missed in the future [.....] I don't know scientifically if it matters but for them it gives them huge reassurance that they've got, it means that rather than them having to see the clinician every week or twice a week, they're doing something, it's rather like feeding a baby every other day."</i> <b>Guys and St Thomas's FG1</b></p> <p><i>"There's a lot of women that have mildly raised blood pressure that are fine, that we don't really do anything with. [Um] and it's a lot of monitoring for them"</i> <b>John Radcliffe, Obstetrician</b></p> <p><i>"some women might not want to monitor themselves because they wouldn't feel, it will be too much responsibility in case they miss something"</i> <b>John Radcliffe FG3</b></p>

	<p><i>"I think it's a bit of mix isn't it, I think some women would really like it saves them time especially if they're busy they have other children."</i> <b>John Radcliffe FG3</b></p> <p><i>"I think a lot of women would also appreciate it because they often don't want to come into hospital and they would, I think if they were self-monitoring at home if they're taught in the right way and they know when somethings high or they know when to escalate or to ring and I think it will work really well and stop a lot of people coming in unnecessarily. I think it would be really good"</i> <b>John Radcliffe, Hospital Midwife</b></p>
<p><b>iv) Anticipated impact on the antenatal care</b></p>	<p><i>"we are so busy, we are too busy in hospitals and the more we can do out of hospital safely the better"</i> <b>John Radcliffe, Obstetrician</b></p> <p><i>"there are benefits all-round aren't there, if you're, if we're [um] happy that this woman is monitoring her blood pressure sensibly [um] it's got benefits for her in that it means she doesn't have to come so often, it's got benefits for us in that it reduces our, our workload, surely it's got, it has a, there's an economic benefit there for the NHS [um] as a whole and it, I suppose that's one of the things isn't it that public health it's about educating people isn't it about, so that, you know, I think there are lots of benefits it's [um] it's just getting it in place. And being confident about the thing that you've got in place"</i> <b>Whipps Cross, Day Assessment Unit midwife</b></p> <p><i>"I think it would have the potential to increase the appointments rather than decrease appointments as a screening tool for hypertension because of the worried well who've seen systolic increase in ten as</i></p>

	<p><i>an example but I don't think it will decrease the routine ante-natal screening because that has many other roles including measurement of fundal height or foetal wellbeing."</i> <b>Royal London, medical trainee</b></p> <p><i>"I suppose potentially it will increase workload because people are going to recognise that they have a slightly elevated blood pressure sooner but I don't think that's necessarily a bad thing"</i> <b>John Radcliffe, Obstetrician</b></p> <p><i>"So it's a massive information imparting exercise probably more importantly than the, the systematic screening for obstetric complications that old fashioned people think of as ante-natal care. You know I, you know, again when I trained we'd see clinics of 40 people, 50 people in a morning where literally all you do would be check their blood pressure, check their urine, put your hand on their tummy, check their heart, you know, see if the patients still breathing, next, you know. But it isn't like that now there's a lot more education, seeing much more as a health education exercise as much as a screening for abnormality of pregnancy exercise".</i> <b>Royal Berkshire, Consultant Obstetrician</b></p>
<p><b>v) Caution, uncertainty and evidence</b></p>	<p><b>Caution</b></p> <p><i>"I wouldn't take the, take everything as a 100% as gospel because, you know, you can't really it's like triaging over the phone they can tell you so much but until you've seen that woman and made an assessment you can't do a full assessment from that. But [um] but yes I mean it would certainly I, I like compare to a jigsaw puzzle, you know, putting all the pieces together that would definitely be one of the pieces I would definitely include once I knew that all the bits were safe that they were using and everything, yes."</i> <b>Royal Wolverhampton FG2</b></p>

*"if I look at it and I can see that yes at 2:00 'o' clock and half two she had raised blood pressure but [...] guidelines about what I do about that [um] all I can do is do my own blood pressure readings and get them reviewed by the doctor. Unless there was like a real, you know, like a guideline of what we actually do about the trends, there's not much I myself could actually do from it."* **Royal**

**Wolverhampton FG4**

#### **Uncertainty**

*"I think that would be very useful information to have rather than as you say the snap shot that you get in clinic. Although on the opposite side if they were having consistently low readings at home and when they come and visit us in triage it's always 160/110 I would question whether the data was true even if I knew that the machine was valid. Acknowledging the role of white coat hypertension but I would still be very worried about relying purely on the home readings and discarding the readings I have in the hospital."* **Royal London, medical trainees.**

*"You can also have high blood pressure at home and lower blood pressure readings in hospital and we have no idea, I don't think [um] what we should do when actually readings are different in different places and there is a tendency, there will be some people who will have a tendency to treat anybody whatever the excuse, much more common is a tendency to say oh it's probably more, we've had 150, 140, 120 I'm going to believe the normal whether its home reading or hospital readings. And so just having more data may just be making more noise rather than giving us, telling us what to do. But without the data we don't even know if that's true."* **Guys and St Thomas's FG1**



	<p><i>"supposed to be evidence based and when you actually look at them about 90% of the recommendations they make are expert opinion"</i> <b>Royal Berkshire, Consultant Obstetrician</b></p>
vi) Concerns over action/inaction	<p><i>"Yes exactly and they may not realise the seriousness of it, so that would be my concern if they're doing it at home. The chronic hypertension women have a lot of contact with, in hypertension clinic they understand blood pressure they've been dealing with it, they're on medication they understand something about blood pressure, they might not fully understand how serious it can be with regards to pre-eclampsia but they certainly understand that they need to act on it and if they don't, obviously it's explained to them numerous times. My concern is if you did that to a general population will that message be translated."</i> <b>Guys and St Thomas's FG2</b></p> <p><i>"I do have a worry about women taking, you know, something like pre-eclampsia is a multi-system disorder, blood pressure is one component and we see every week, every month women who come in with normal blood pressure but everything else going wrong and it would make me nervous about the idea of women taking control of their care so much so that they felt reassured by one reading and ignored their signs and symptoms and I think that, that will be, that's a major, that's a clinical worry that I have for using."</i> <b>John Radcliffe, Consultant Obstetrician</b></p> <p><i>"I think as long as they [women] understand the limits of what's normal and what's not and they know the right people to contact if not, then I don't see there's anything wrong with it"</i> <b>Royal Berkshire FG2</b></p>

*"I just know that for some women they, they've got so much going on that actually their health is, is quite easily overlooked and [um] is last on the agenda if that makes sense and if they could miss a blood pressure reading here and there [um] then they may well do that and that would be my only [um] and which I feel awful saying really because I am so like passionate about actually women being responsible for their own care and taking ownership of it and us giving it back to them [um]. [.....] I just think some women will be like oh yes, everything's fine they're all okay and maybe overlook things. I know you've got the opposite extent where some people would potentially act on it like say there's a higher reading because they think it will be acted on and that sort of thing as well but I think you could almost get women who will either pretend they've done it or pretend that it's an okay reading when it's not [um] which is awful to say that we don't trust women but"* **Royal Wolverhampton FG2**

# Reporting checklist for qualitative study.

Based on the SRQR guidelines.

## Instructions to authors

Complete this checklist by entering the page numbers from your manuscript where readers will find each of the items listed below.

Your article may not currently address all the items on the checklist. Please modify your text to include the missing information. If you are certain that an item does not apply, please write "n/a" and provide a short explanation.

Upload your completed checklist as an extra file when you submit to a journal.

In your methods section, say that you used the SRQRreporting guidelines, and cite them as:

O'Brien BC, Harris IB, Beckman TJ, Reed DA, Cook DA. Standards for reporting qualitative research: a synthesis of recommendations. Acad Med. 2014;89(9):1245-1251.

		Page Number
Reporting Item		
Title		
#1	Concise description of the nature and topic of the study identifying the study as qualitative or indicating the approach (e.g. ethnography, grounded theory) or data collection methods (e.g. interview, focus group) is recommended	1
Abstract		
#2	Summary of the key elements of the study using the abstract format of the intended publication; typically includes background, purpose, methods, results and conclusions	2
Introduction		
Problem formulation	#3 Description and significance of the problem / phenomenon studied: review of relevant theory and empirical work; problem statement	4
Purpose or research question	#4 Purpose of the study and specific objectives or questions	5

## Methods

Qualitative approach and research paradigm	<a href="#">#5</a>	Qualitative approach (e.g. ethnography, grounded theory, case study, phenomenology, narrative research) and guiding theory if appropriate; identifying the research paradigm (e.g. postpositivist, constructivist / interpretivist) is also recommended; rationale. The rationale should briefly discuss the justification for choosing that theory, approach, method or technique rather than other options available; the assumptions and limitations implicit in those choices and how those choices influence study conclusions and transferability. As appropriate the rationale for several items might be discussed together.	5
Researcher characteristics and reflexivity	<a href="#">#6</a>	Researchers' characteristics that may influence the research, including personal attributes, qualifications / experience, relationship with participants, assumptions and / or presuppositions; potential or actual interaction between researchers' characteristics and the research questions, approach, methods, results and / or transferability	5
Context	<a href="#">#7</a>	Setting / site and salient contextual factors; rationale	5
Sampling strategy	<a href="#">#8</a>	How and why research participants, documents, or events were selected; criteria for deciding when no further sampling was necessary (e.g. sampling saturation); rationale	5
Ethical issues pertaining to human subjects	<a href="#">#9</a>	Documentation of approval by an appropriate ethics review board and participant consent, or explanation for lack thereof; other confidentiality and data security issues	16
Data collection methods	<a href="#">#10</a>	Types of data collected; details of data collection procedures including (as appropriate) start and stop dates of data collection and analysis, iterative process, triangulation of sources / methods, and modification of procedures in response to evolving study findings; rationale	5
Data collection instruments and technologies	<a href="#">#11</a>	Description of instruments (e.g. interview guides, questionnaires) and devices (e.g. audio recorders) used for data collection; if / how the instruments(s) changed over the course of the study	5

1	Units of study	<a href="#">#12</a>	Number and relevant characteristics of participants, documents, or events included in the study; level of participation (could be reported in results)	5
2				
3				
4				
5				
6	Data processing	<a href="#">#13</a>	Methods for processing data prior to and during analysis, including transcription, data entry, data management and security, verification of data integrity, data coding, and anonymisation / deidentification of excerpts	6
7				
8				
9				
10				
11				
12				
13	Data analysis	<a href="#">#14</a>	Process by which inferences, themes, etc. were identified and developed, including the researchers involved in data analysis; usually references a specific paradigm or approach; rationale	6
14				
15				
16				
17				
18	Techniques to enhance trustworthiness	<a href="#">#15</a>	Techniques to enhance trustworthiness and credibility of data analysis (e.g. member checking, audit trail, triangulation); rationale	6
19				
20				
21				
22				
23				
24	<b>Results/findings</b>			
25				
26	Syntheses and interpretation	<a href="#">#16</a>	Main findings (e.g. interpretations, inferences, and themes); might include development of a theory or model, or integration with prior research or theory	6
27				
28				
29				
30				
31	Links to empirical data	<a href="#">#17</a>	Evidence (e.g. quotes, field notes, text excerpts, photographs) to substantiate analytic findings	7-11, Table 2
32				
33				
34				
35	<b>Discussion</b>			
36				
37				
38	Intergration with prior work, implications, transferability and contribution(s) to the field	<a href="#">#18</a>	Short summary of main findings; explanation of how findings and conclusions connect to, support, elaborate on, or challenge conclusions of earlier scholarship; discussion of scope of application / generalizability; identification of unique contributions(s) to scholarship in a discipline or field	12-14
39				
40				
41				
42				
43				
44				
45				
46	Limitations	<a href="#">#19</a>	Trustworthiness and limitations of findings	13
47				
48	<b>Other</b>			
49				
50				
51	Conflicts of interest	<a href="#">#20</a>	Potential sources of influence of perceived influence on study conduct and conclusions; how these were managed	15
52				
53				
54	Funding	<a href="#">#21</a>	Sources of funding and other support; role of funders in data collection, interpretation and reporting	16
55				
56				
57				
58	Notes:			
59				
60				

- 17: 7-11, Table 2 The SRQR checklist is distributed with permission of Wolters Kluwer © 2014 by the Association of American Medical Colleges. This checklist was completed on 19. February 2020 using <https://www.goodreports.org/>, a tool made by the [EQUATOR Network](#) in collaboration with [Penelope.ai](#)

For peer review only

# BMJ Open

## Exploring the potential for introducing home monitoring of blood pressure during pregnancy into maternity care: current views and experiences of staff. A qualitative study.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2020-037874.R1
Article Type:	Original research
Date Submitted by the Author:	21-Jul-2020
Complete List of Authors:	Hinton, Lisa; University of Cambridge, THIS Institute, Department of Public Health and Primary Care Hodgkinson, James ; University of Birmingham, Primary Care Clinical Sciences, Institute of Applied Health Research Tucker, Katherine; University of Oxford, Nuffield Department of Primary Care Rozmovits, Linda; Freelance Researcher Chappell, Dr Lucy; King's College London, Women's Health Academic Centre Greenfield, Sheila; University of Birmingham, Primary Care Clinical Sciences McCourt, Christine; City University of London, Department of Midwifery and Child Health Sandall, Jane; Kings College, London, Health and Social Care Research Division McManus, Richard; University of Oxford, Dept of Primary Care Health Sciences
<b>Primary Subject Heading</b>:	Obstetrics and gynaecology
Secondary Subject Heading:	Health services research
Keywords:	Hypertension < CARDIOLOGY, Organisation of health services < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Maternal medicine < OBSTETRICS, QUALITATIVE RESEARCH

SCHOLARONE™  
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.



**Exploring the potential for introducing home monitoring of blood pressure during pregnancy into maternity care: current views and experiences of staff. A qualitative study.**

Hinton L<sup>1</sup>, Hodgkinson J<sup>3</sup>, Tucker K<sup>2</sup>, Rozmovits L<sup>5</sup>, Chappell LC<sup>4</sup>, Greenfield S<sup>3</sup>, McCourt C<sup>6</sup>, Sandall J<sup>4</sup>, McManus RJ<sup>2</sup>

<sup>1</sup> THIS Institute, Department of Public Health and Primary Care, University of Cambridge, CB2 0AH.

<sup>2</sup> Nuffield Department of Primary Care Health Sciences, University of Oxford OX2 6GG, UK.

<sup>3</sup> Institute of Applied Health Research, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK.

<sup>4</sup> Department of Women and Children's Health Academic Centre, King's College London, London, SE1 7EH

<sup>5</sup> Freelance Researcher, Montreal, Quebec H2L 3X1, Canada

<sup>6</sup> Centre for Maternal and Child Health, School of Health Sciences, City University, London EC1R 1UW, UK

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Running title:** Staff views on home BP monitoring during pregnancy

**Author responsible for correspondence:** Dr Lisa Hinton

Tel +44 (0)1223 731573    lisa.hinton@thisinstitute.cam.ac.uk

THIS Institute, Department of Public Health and Primary Care, University of Cambridge.

Cambridge Biomedical Campus, Cambridge CB2 0AH.

For peer review only

## ABSTRACT

**Objective** One in 20 women are affected by pre-eclampsia, a major cause of maternal and perinatal morbidity, death and premature birth worldwide. Diagnosis is made from monitoring blood pressure (BP) and urine and symptoms at antenatal visits after 20 weeks of pregnancy. There are no randomized data from contemporary trials to guide the efficacy of self-monitoring of blood pressure (SMBP) in pregnancy. We explored the perspectives of maternity staff to understand the context and health system challenges to introducing and implementing SMBP in maternity care, ahead of undertaking a trial.

**Design.** Exploratory study using a qualitative approach.

**Setting.** Eight hospitals, English National Health Service.

**Participants** Obstetricians, community and hospital midwives, pharmacists, trainee doctors (n=147)

**Methods** Semi-structured interviews with site research team members and clinicians, interviews and focus group discussions. Rapid content and thematic analysis undertaken.

**Results** The main themes to emerge around SMBP include i) different blood pressure changes in pregnancy ii) reliability and accuracy of BP monitoring iii) anticipated impact of SMBP on women iv) anticipated impact of SMBP on the antenatal care system, v) caution, uncertainty and evidence vi) concerns over action/inaction and patient safety.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Conclusions** The potential impact of SMBP on maternity services is profound although  
nuanced. While introducing SMBP does not reduce the responsibility clinicians have for  
women’s health, it may enhance the responsibilities and agency of pregnant women, and  
introduces a new set of relationships into maternity care. This is a new space for  
reconfiguration of roles, mutual expectations, and the relationships between and  
responsibilities of health care providers and women.

### Strengths and limitations of this study

- Qualitative study to explore maternity staff perspectives on self-monitoring of blood pressure in pregnancy
- Diverse sample includes voices from across multidisciplinary maternity teams
- Focus groups held across 8 hospitals in the English National Health Services, including hospital and community midwives

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**MAIN BODY OF TEXT**

Introduction

Raised blood pressure (BP) in pregnancy affects around one in ten women, almost half of whom develop pre-eclampsia, a major cause of maternal and perinatal morbidity, death and premature birth worldwide.(1, 2) Diagnosis is made from monitoring BP and urine and symptoms at antenatal visits after 20 weeks of pregnancy.

Self-monitoring of BP (SMBP) in the general population has become more common and large trials have demonstrated its effectiveness in terms of BP control. (3) Since 2011 UK national guidance for the general population with hypertension there has been a growing acceptance by clinicians of using SMBP values. Primary care physicians report an increasing use of SMBP to diagnose hypertension and for the ongoing management of hypertension to monitor BP control. (4-7)

But what are the implications of introducing SMBP for women during or after pregnancy, to improve the detection and management of gestational hypertension and pre-eclampsia? Pregnant women are different from the general population: BP can rise rapidly during pregnancy, and the problems associated with this (such as pre-eclampsia, eclampsia and

stroke) are potentially serious. Pilot work in the UK suggests, with support from midwives and doctors, it is feasible and acceptable for women to monitor their BP and urine safely, potentially identifying hypertension earlier and controlling BP better.(8-10) SMBP during pregnancy is becoming common in some settings. A Canadian pilot study found more than 60% of women diagnosed with non-proteinuric hypertension in pregnancy were already undertaking SMBP. (11) The Society of Obstetricians and Gynaecologists of Canada provides guidelines for SMBP in pregnancy and most Canadian obstetricians and primary care physicians used SMBP to assess for white coat hypertension (WCH), (12) but its place in other countries is not yet clear. (13)

While SMBP in pregnancy is already implemented in some UK settings, (14) there are no adequately powered randomized data from contemporary trials to guide the efficacy of such methods. Our early work exploring women's experiences of taking part in an SMBP pilot study (15) showed that most found their health professionals to be supportive, although in some cases self-monitoring results were ignored. An understanding of the attitudes of individuals and the organization will be paramount in introducing new, and safe, models of care. Ahead of undertaking a large trial of BP monitoring in high-risk pregnancy to improve detection and monitoring of hypertension, we undertook a study to explore the perspectives of staff across

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

the maternity care pathway to explore the context and health system challenges to introducing and implementing SMBP in maternity care. (10, 16, 17)

Methods

We carried out an intervention development phase in preparation for a large trial of SMBP in pregnancy, using qualitative methods with the aim of understanding the views and prior experience of clinicians regarding such monitoring. This involved interviews and focus groups across eight participating teaching and urban general district hospitals that serve diverse populations. (TABLE 1). We conducted face-to-face interviews with a lead consultant obstetrician at each site (n=8) and a series of focus groups and interviews with a wide range of healthcare professionals.

**Sample/setting** We sought a broad pragmatic sample that included community and hospital-based midwives and obstetricians of different seniorities at each site (n=147). All focus groups and interviews took place at hospital or health centre sites, either as part of, or after, existing meetings or during shift breaks. It was usually a Research Midwife at each site who identified who might be available with some of the larger focus groups piggy-backing on existing meetings. Sampling was therefore convenience rather than purposive, though we ensured there was a representation of senior and junior obstetricians, and hospital and community midwives from each site. Most participants had not previously engaged in



previous studies on SMBP. Focus group size ranged from two to 15 individuals. See Table

1. Some focus groups had a mix of obstetricians and midwives participating, reflecting the

multidisciplinary teams providing care to women during pregnancy. Although an obstetrician

is the lead clinician for almost all women with additional risk factors in pregnancy, the

midwife is the conduit of care throughout pregnancy, labour and the postnatal period. (18)

TABLE 1 Study Sample

HOSPITAL	FOCUS GROUP	NUMBERS	Staff represented
Guys & St Thomas's NHS Foundation Trust	Focus group 1	10	Obstetricians, midwives, pharmacist
	Focus group 2	4	Day assessment unit midwives
	PI Interview	1	Consultant Obstetrician
West Middlesex University Hospital	Focus group 1	3	Midwives
	Focus group 2	4	Midwives
	Focus group 3	5	Midwives and obstetrician
	Focus group 4	3	Midwives
	PI Interview	1	Consultant Obstetrician
Whipps Cross Hospital, Barts Health NHS Trust	Focus group 1	11	Obstetricians, midwives
	Focus group 2	4	Antenatal midwives
	Focus group 3	4	Day assessment unit midwives
	PI Interview	1	Consultant Obstetrician
The Royal London Hospital, Barts Health NHS Trust	Focus group 1	6	Community midwives (FMU)
	Focus group 2	4	Community midwives (in hospital)
	Interview	1	Consultant obstetrician
	Focus group 3	7	Antenatal midwives
	Focus group 4	2	Medical trainees

	PI Interview	1	Consultant Obstetrician
John Radcliffe Hospital, Oxford University Hospitals NHS Foundation Trust	Focus group 1	11	Community midwives (in hospital)
	Focus group 2	3	Obstetricians
	Interview	1	Obstetrician
	Interview	1	Consultant obstetrician
	Interview	1	Obstetrician
	Focus Group 3	2	Hospital midwives (MAU)
	Interview	1	Hospital midwife (MAU)
	PI Interview	1	Consultant Obstetrician
Horton General Hospital, Oxford University Hospitals NHS Foundation Trust	Focus Group 4	13	Community midwives
Royal Berkshire NHS Foundation Trust	Focus group 1	2	Hospital midwives
	Focus group 2	2	Hospital midwives
	Interview	1	Hospital midwife
	Focus group 3	2	Obstetricians
	Interview	1	Obstetrician
	Focus group 4	6	Community midwives (in hospital)
	PI Interview	1	Consultant Obstetrician
The Royal Wolverhampton NHS Trust	Focus group 1	3	Obstetricians
	Focus group 2	2	Hospital midwives
	Focus group 3	10	Hospital and community midwives

	Focus group 4	8	Hospital and community midwives
	Interview	1	Hospital midwife
	PI Interview	1	Consultant Obstetrician
Birmingham Women's NHS Foundation Trust	PI Interview	1	Consultant Obstetrician
	<b>Total</b>	<b>147</b>	

**Data collection** The interviews and focus groups were facilitated by two experienced, post-doctoral social science/non-clinical researchers (LH and JH, male and female) and audio-recorded for transcription and analysis. A topic guide was developed drawing on the literature, our prior work and in discussion with the multi-disciplinary trial team (see Topic Guide, Supplementary file). Questions sought to explore issues around staff views on SMBP in pregnancy, the accuracy of home readings, how to manage care of women who present with home readings (particularly those discordant with clinic readings), the impact of home readings on clinical decision-making and on workload, as well as practical issues around operationalising and integrating self-monitoring into current care pathways. Ethical approval and informed consent was obtained.

**Analysis** All interviews and discussions were transcribed for analysis, which was conducted in two phases. An initial rapid analysis was undertaken to guide trial development. (19) This used templates developed by JH and LH based on the topic guides and subsequently refined after

1  
2  
3 a period of piloting. We conducted a rapid content analysis to identify evidence related to the  
4  
5  
6 a priori concepts and emergent themes that would then guide development of the study. This  
7  
8  
9 was discussed and refined with the wider research team. In a subsequent phase LH and JH,  
10  
11  
12 in discussion with the wider team, developed a coding frame and worked in parallel (with LR)  
13  
14  
15 to code and analyse the data.(20) Thematic analysis was undertaken to explore the  
16  
17  
18 anticipated and emergent themes and was then mapped back to the framework, supported by  
19  
20  
21 NVIVO (21, 22). LH and JH met frequently to discuss the results together and then with the  
22  
23  
24 wider team to confirm the credibility and dependability of the analysis. Data collection  
25  
26  
27 continued until data saturation was reached for the themes reported in this paper. (23)  
28  
29  
30 Interviews and focus groups were analysed as a single dataset, reflecting the multidisciplinary  
31  
32  
33 teams providing care to women with additional risk factors in pregnancy. Results are  
34  
35  
36 presented as staff views, with illustrative quotes presented in TABLE 2.  
37  
38  
39

40  
41 **Patient and public involvement** This paper reports a sub-study of a programme of intervention  
42  
43  
44 development work for the BUMP trials.(24) There was no direct patient involvement in the  
45  
46  
47 research reported here, as the focus was staff views and experiences. However patient  
48  
49  
50 priorities were central to developing the trials and intervention development work reported  
51  
52  
53 elsewhere. (25)  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

RESULTS

We heard views from a total of 147 healthcare professionals providing maternity care in the English NHS, of whom 37 were physicians (including PIs and students) 109 midwives and one pharmacist. See Table 1. Our findings suggest SMBP during pregnancy was not yet a widespread practice across the sites included. However, some midwives and obstetricians reported occasional pregnant women who were already self-monitoring, either of their own volition or because they had been advised to because of previous experiences of pre-eclampsia, white coat hypertension (WCH), or pre-existing hypertension. The main themes to emerge around SMBP were i) different blood pressure changes in pregnancy ii) reliability and accuracy of BP monitoring iii) anticipated impact of SMBP on women iv) anticipated impact of SMBP on the antenatal care system, v) caution, uncertainty and evidence vi) concerns over action/inaction and patient safety.

*Interpreting blood pressure changes in pregnancy*

BP fluctuation

Staff emphasised that BP is different in pregnancy than at other times in women’s lives. It fluctuates throughout pregnancy and these fluctuations are normal. However, BP in pregnancy can change rapidly, precipitating a medical emergency for mother and baby. While antenatal care staff will be familiar with this phenomenon through their training and

1  
2  
3 experience, women may not be. There was concern women could have difficulty interpreting  
4  
5  
6 results and deciding when to seek acute care.  
7  
8  
9

10 In the general population, BP readings are used to guide care and medication changes over  
11  
12 the long term, but in pregnancy the situation is more dynamic. Some were interested in the  
13  
14 wider context home BP readings would give them, in particular, feeling women with WCH  
15  
16 might get more typical readings at home. However, several indicated they would tend to  
17  
18 privilege their own clinic readings as they reflected the situation happening immediately in  
19  
20 front of them.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

### 32 Wider symptoms

33 Staff were also concerned that BP is not the only factor in diagnosing hypertension or pre-  
34  
35 eclampsia but rather one piece of a diagnostic 'jigsaw'. Healthcare professionals stressed the  
36  
37 importance of ensuring women were alert to the wide range of symptoms (e.g. headaches,  
38  
39 dizziness, blurred vision, proteinuria) and would act on those even if their BP appeared to be  
40  
41 in the normal range.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

### 53 *Reliability and accuracy*

54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The reliability and accuracy of machines were key concerns for staff, as well as whether women would monitor their BP correctly and respond appropriately. Many raised concerns about the quality of home BP monitors, whether women were using one of the relatively few validated for use in pregnancy, and whether they were properly calibrated. There were concerns about appropriate cuff size, particularly for women with a high body mass index. Other factors identified as potentially impacting the reliability and accuracy of home readings included women’s practices and the context in which they took readings (did they sit correctly, choose a quiet time of day?).

*Views on the impact on women*

Staff had a range of perceptions about how pregnant women might feel about being asked to self-monitor their BP. Some anticipated women being happy to participate and enjoying a sense of agency, control and reassurance over their own healthcare. Others raised concern that women may experience increased anxiety and become obsessed about their BP. Some felt pregnant women expect care to be led by providers and asking them to monitor their own BP would be perceived as a way of abdicating responsibility. Although staff also acknowledged the potential for empowerment, some expressed concern that asking women



1  
2  
3 to self-monitor would contribute to an over-medicalisation of pregnancy in contrast with the  
4  
5  
6 evolution of antenatal care away from this model in recent years.  
7  
8  
9

10  
11  
12  
13 Many potentially positive impacts for women were identified both in terms of quality of  
14  
15  
16 experience and better outcomes for mother and baby. These included fewer medical  
17  
18  
19 appointments, shorter waiting times and avoiding hospitalisation. Staff acknowledged  
20  
21  
22 detecting BP increases earlier would have the potential to deliver better outcomes for both  
23  
24  
25 mothers and babies, but confirmed definitive evidence of these outcomes would be needed  
26  
27  
28 before they could commit to changing their practice.  
29  
30  
31  
32  
33  
34

### 35 *Anticipated impact on antenatal care*

36  
37  
38  
39

40 Staff were caught between feeling SMBP could reassure women, so cutting 'worried well' visits  
41  
42  
43 and/or raise anxieties, thus increasing them . So, while they recognised other potential health  
44  
45  
46 system benefits including fewer women coming into clinic, thus freeing up clinic time in hospital  
47  
48  
49 and in community care, they also raised various concerns around the impact of SMBP on  
50  
51  
52 antenatal care pathways and their workload. These included a rise in phone calls to maternity  
53  
54  
55 assessment units and an increased demand for care, particularly from the 'worried well'.  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Staff were aware that, ahead of definitive trial results, these reflections were hypothetical.

Their currently available best comparator was their experience with women’s self-monitoring for gestational diabetes, which is now quite common. Some described SMBP as a challenge to their professional roles, particularly for midwives, and part of wider changes towards increasing self-care and personal responsibility in health. One consultant described the shifting responsibilities in antenatal care, with health education now as much a part of routine care as systematic screening for obstetric complications.

*Caution, uncertainty and evidence*

While staff were generally open to looking at home BP readings, most were mindful the evidence was not yet available as to how to treat these readings in conjunction with clinic readings. Most felt they could only treat home readings as additional rather than as core information, not directing care decisions, but contributing extra detail to the ‘jigsaw’:

**Caution**

Many midwives said that if self-monitoring readings were discrepant with clinic readings, they would privilege the readings they had just taken themselves. Most added they would take into account the home readings, though exactly how would be considered on a case-by-case

1  
2  
3 basis. Some would use them as part of improving understanding of the wider context, but use  
4  
5  
6 their clinic reading for the decision on whether to admit a woman to hospital. The home reading  
7  
8  
9 would not be used for treatment decisions but could influence how often an obstetrician felt it  
10  
11  
12 necessary to see a woman.  
13  
14  
15

16  
17  
18  
19 Some acknowledged the potential of SMBP to identify WCH. But in several midwife  
20  
21  
22 discussions, consensus emerged on the side of caution, such as referring women onwards or  
23  
24  
25 taking bloods if only one type of reading (clinic or home) was high. Others referred to local  
26  
27  
28 protocols and guidelines although this was often interpreted as needing to act on any higher  
29  
30  
31 reading, implying caution and a focus on patient safety, with fear of being considered negligent  
32  
33  
34 explicitly raised by one midwife.  
35  
36  
37

38  
39  
40  
41 Some said they verified home readings by testing the device used by women alongside their  
42  
43  
44 clinic monitor. Several midwives commented they would have to respect a woman's readings,  
45  
46  
47 as to do otherwise could affect mutual trust.  
48  
49  
50

## 51 52 53 54 Uncertainty

55  
56 There was awareness that pre-eclampsia is a variable condition in which the speed of disease  
57  
58  
59 progression varies considerably. Some felt SMBP could significantly mitigate some of these  
60

1  
2  
3 uncertainties. One senior obstetrician specifically alluded to a potential *advantage* of SMBP  
4  
5  
6 being its ability to reduce uncertainty by enabling clinicians to make objective assessment of  
7  
8  
9 how accurate clinic readings are, clarifying who has WCH, assessing the effectiveness of  
10  
11  
12 treatment and standardising the various clinical environments that women might be seen in.  
13  
14  
15  
16 Yet others clearly feared SMBP could increase uncertainty (discrepancy in readings), and/or  
17  
18  
19 lead to overcaution and overtreatment on the part of less experienced colleagues.  
20  
21  
22  
23  
24  
25

26 Different approaches to managing hypertension in pregnancy also emerged. While some  
27  
28  
29 make judgements on a case-by-case basis, privileging the extent of change from baseline, the  
30  
31  
32 wider context, and individualised pathways, and given that different hospitals may have their  
33  
34  
35 own guidelines, others described a practice defined by the National Institute for Health and  
36  
37  
38 Care Excellence (NICE) guidelines and the use of thresholds as the defining guide in when to  
39  
40  
41 intervene (26).  
42  
43  
44  
45  
46  
47

48 Evidence  
49  
50 Numerous staff raised the need for more and improved evidence. One obstetrician observed  
51  
52  
53 that guidelines are supposed to be evidence based but recommendations are often made on  
54  
55  
56 expert opinion. A different senior obstetrician also suggested that people practise defensively,  
57  
58  
59 leading to many unnecessary blood tests being done because a greater number of healthcare  
60

professionals (midwives and junior doctors) are able to request the tests, and because when women have been told they may have pre-eclampsia, heightened anxiety leads to repeated checks for reassurance. Overall, there was a distinct lack of consensus, but with a clear wish from many for more evidence to reduce uncertainty.

### *Concerns over action/inaction*

While a shift to SMBP is in keeping with wider trends towards self-care and personal responsibility, working through this new balance of responsibility for acting on readings was a cause for concern. Staff raised concerns about women not acting on readings, through either not realising they required urgent action, not wanting to bother busy staff, or not wanting to be hospitalised. There were concerns about women ignoring problems, or being falsely reassured by one reading and ignoring the wider context. Many stressed the importance of women having a clear understanding of what they were supposed to do if their readings were high, with a clear pathway for women presenting with high BP based on home readings.

Many staff made comparisons with home monitoring of blood glucose in gestational diabetes, either as a useful precedent for SMBP or evidence of the challenges of women deliberately or inadvertently coming in with inaccurate results.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Others expressed the view that it is hospital policy (as well as being the health professional's view) that they would rather ask women to come in for comprehensive assessment than have the woman check her BP at home and decide not to come in.

DISCUSSION

Main findings.

Antenatal care staff were generally in favour of asking women to monitor their own BP at home, but flagged concerns about the potential impacts on women and staff workload, the reliability and accuracy of readings and the need for clarity about how the information would be used by clinicians. Many emphasised the importance of ensuring women understood what to do in the event of readings that cause concern and provided examples of other situations where women had failed to act in acute situations because they did not understand the severity of the situation or rationalised away the need to seek medical attention. Some voiced concerns about women becoming either anxious or falsely reassured by self-monitoring. There was a feeling that SMBP might only be for a select few as it will require a shift in thinking about who takes responsibility for the woman's health.

Many of the issues raised were specific to pregnancy. However, concerns about the potential for monitoring leading to anxiety may also exist for the general population – though previous studies have not found evidence to support this (27) – they could be intensified when there may be additional anxiety related to the effect of BP changes on the baby. Concerns about device accuracy were well founded as relatively few monitors on the market are validated in pregnancy.(28) The need to consider more than just BP and the dynamic nature of BP in pregnancy as opposed to BP in the general population means that failure to act has the potential for devastating consequences for both woman and baby. Yet resistance to medicalisation may be heightened in a previously healthy younger population of ‘women’ as opposed to ‘patients’. (29) Continuity of care is an ambition not always achieved in maternity care. The lack of an established relationship of trust between women/patients and healthcare professionals that is typical of hypertension management in primary care, further complicates the situation (30). Staff are therefore understandably cautious and uncertain, while also willing to see the potential for SMBP to reduce uncertainty and add usefully to available evidence.

## Strengths and limitations

A strength of this study is the breadth of multidisciplinary voices across the maternity care pathway captured, spanning eight differently located hospitals in England. This has given rich insights into the potential challenges of introducing SMBP into maternity care from the

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

perspectives of different disciplines and locales. Our two-stage approach to analysis, led by social scientists with expert clinical input from co-authors in obstetrics, midwifery and primary care, has also enhanced the interpretive strength of the findings. Limitations include the challenges of holding focus group discussions with busy professionals engaged with frontline care. These are vital perspectives to capture but there were inevitably challenges in terms of finding time and space for some focus group discussions.

There are several literatures to draw on in interpreting these results.

**Responsibilisation:**To date, measuring and monitoring women’s BP in pregnancy has been a central component of the healthcare professional’s role in antenatal care. To include women’s home readings into these care pathways represents a significant shift in the balance of responsibility. Emerging literature that explores the shifts and implications of health ‘responsibilisation’ are of relevance. Shifts in the balance of responsibility have the potential to disrupt long established care pathways and can be experienced both positively as well as negatively as surveillance.(31) Exploring the impacts of these shifts is a live issue in the sociological literature from studies of ‘park runs’ to HIV management (32, 33), and the relationships between social workers and vulnerable youth .(34) Drawing on work by Garret (2008), Liebenberg et al explored the impact of shifts in neo-liberal systems, where individuals



are expected to manage their own risks and demonstrate self-care. “*These shifts impact on the roles that health professionals play, from active management to encouraging/teaching of the skills of management*” (35). Newman et al’s 2012 study of young people with HIV revealed narratives of responsibilisation that can give rise to contest between young people and their clinicians. (33) In exploring the impact of telecare on the management of long term conditions, Rogers et al (2011) concluded “*Indeed a paradox of the reliance and acceptance of telecare is the creation of new relationships and dependencies rather than the diminution of reliance envisaged by policy.*” (36) Our findings on clinician caution and concerns over inaction, indicate that these shifts in responsibility could have profound impacts that we need to be alive to as we further explore the impact of this work going forward. (37)

**Impact on professional roles:** A shift in responsibility for BP monitoring may also have an impact on professional roles and identity. Concern over these impacts ran through various themes we have reported, including impact on workload, caution and uncertainty and location of expertise. Task shifting is a key issue in many healthcare settings, but it does not emerge seamlessly as revealed in the literature.(38-41) How SMBP impacts on these professional roles, and how professionals behave when there is a shift in the locus of responsibility will need to be the focus for future work.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Patient safety:** While staff raised concerns about the impact of SMBP on patient safety, these concerns were expressed in the context of an emerging evidence base for SMBP (13, 17). As this develops, we draw on evidence that points to the considerable scope for improving patient and family contributions to the detection and management of acute illness (42). Delayed recognition and treatment of conditions such as pneumonia and meningitis in childhood (43), pre-eclampsia and reduced fetal movements during pregnancy and after childbirth (44, 45) and heart disease and stroke in adulthood, (46, 47) contribute significantly to the mortality and morbidity burden. These conditions typically present with a time-critical window for early recognition and response, and are associated with red flag signs and symptoms (such as breathlessness and pain) which can signify a serious underlying condition and act as potential markers to aid patient and family involvement in escalation of care. However, research has shown there are challenges to speaking up and raising safety alerts in maternity care and that organization-focused efforts are required to improve staff responsiveness. (48)

## Conclusions

We need to be mindful that the impact of SMBP on maternity services could be profound although nuanced. Whilst its introduction will not reduce the responsibility clinicians have for women's health, it promises to enhance the responsibilities and agency of women, and introduces a new set of relationships into maternity care. We cannot simply look to the literature and experiences of SMBP in other populations (e.g. such as hypertension outside of pregnancy) because of the very nature of BP changes in pregnancy and because this is a healthy population, unless specifically indicated otherwise. This is, then, a new space for reconfiguration of roles, mutual expectations, and the relationships between, and responsibilities of, health care providers and mothers.

## ACKNOWLEDGEMENTS

We would like to thank all NHS staff who gave their time to contribute to this study.

## DISCLOSURE OF INTERESTS

RJM has received BP monitors for research from Omron. No other authors have conflicts to declare.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

CONTRIBUTION OF AUTHORSHIP

Authors contributions are as follows: LH, JH, KT, LCC, SG, CM, JS, LR and RM conceived the study, secured the funding and were involved in the planning and carrying out the study. LH and JH conducted the interviews and focus groups and KT managed the study. LH, JH and LR analysed the data with input from the wider authorship. LH and JH drafted the paper. All authors commented on drafts and writing up the work.

DETAILS OF ETHICS APPROVAL

A favourable ethical review for this study was obtained from South Central - Oxford C Research Ethics Committee (16/SC/0386). Written informed consent was gained from individuals who agreed to take part.

FUNDING

This article represents independent research commissioned by the National Institute for Health Research (NIHR) Programme for Applied Research (RP-PG-0614-20005). Richard McManus was and Lucy Chappell is supported by a Research Professorship from the National Institute for Health Research (NIHR-RP-R2-12-015 and RP-2014-05-019 respectively). Richard McManus in an NIHR Senior Investigator. Lisa Hinton, Katherine Tucker and Richard McManus have received funding from the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care Oxford at Oxford Health NHS Foundation Trust and NIHR Oxford Thames Valley Applied Research Collaboration. Lisa Hinton is also supported by the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC). Jane Sandall is a National Institute for

1  
2  
3 Health Research (NIHR) Senior Investigator and is supported by the National Institute for  
4 Health Research (NIHR) Applied Research Collaboration South London (NIHR ARC South  
5 London) at King's College Hospital NHS Foundation Trust. The views expressed in this  
6 publication are those of the author(s) and not necessarily those of the NHS, the National  
7 Institute for Health Research or the Department of Health.  
8  
9  
10  
11  
12

13  
14 **DATA AVAILABILITY** The interview transcripts are held by the University of Oxford and  
15  
16 available at reasonable request.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

REFERENCES

1. Mol BW, Roberts CT, Thangaratinam S, Magee LA, De Groot CJ, Hofmeyr GJ. Pre-eclampsia. *The Lancet*. 2016;387(10022):999-1011.

2. Conti-Ramsden F, Knight M, Green M, Shennan AH, Chappell LC. Reducing maternal deaths from hypertensive disorders: learning from confidential inquiries. *Bmj*. 2019;364:l230.

3. McManus R, Wood S, Bray E, Glasziou P, Hayen A, Heneghan C, et al. Self-monitoring in hypertension: a web-based survey of primary care physicians. *Journal of human hypertension*. 2014;28(2):123.

4. National Clinical Guideline Centre. Hypertension: management of hypertension in adults in primary care. London: National Institute for Clinical Excellence; 2011.

5. McManus RJ, Mant J, Bray EP, Holder R, Jones MI, Greenfield S, et al. Telemonitoring and self-management in the control of hypertension (TASMINH2): a randomised controlled trial. *The Lancet*. 2010;376(9736):163-72.

6. McManus RJ, Mant J, Franssen M, Nickless A, Schwartz C, Hodgkinson J, et al. Efficacy of self-monitored blood pressure, with or without telemonitoring, for titration of antihypertensive medication (TASMINH4): an unmasked randomised controlled trial. *The Lancet*. 2018;391(10124):949-59.

7. Fletcher BR, Hinton L, Bray EP, Hayen A, Hobbs FR, Mant J, et al. Self-monitoring blood pressure in patients with hypertension: an internet-based survey of UK GPs. *Br J Gen Pract*. 2016;66(652):e831-e7.

8. Tucker KL, Taylor KS, Crawford C, Hodgkinson JA, Bankhead C, Carver T, et al. Blood pressure self-monitoring in pregnancy: examining feasibility in a prospective cohort study. *BMC pregnancy and childbirth*. 2017;17(1):442.

9. Tucker KL, Bowen L, Crawford C, Mallon P, Hinton L, Lee M-M, et al. The feasibility and acceptability of self-testing for proteinuria during pregnancy: A mixed methods approach. *Pregnancy hypertension*. 2018;12:161-8.

10. Mackintosh N, Rance S, Carter W, Sandall J. Working for patient safety: a qualitative study of women's help-seeking during acute perinatal events. *BMC pregnancy and childbirth*. 2017;17(1):232.

11. Magee LA, von Dadelszen P, Chan S, Gafni A, Gruslin A, Helewa M, et al. Women's views of their experiences in the CHIPS (Control of Hypertension in Pregnancy Study) Pilot Trial. *Hypertension in Pregnancy*. 2007;26(4):371-87.

12. Dehaeck U, Thurston J, Gibson P, Stephanson K, Ross S. Blood pressure measurement for hypertension in pregnancy. *Journal of Obstetrics & Gynaecology Canada: JOGC*. 2010;32(4):328-34.

13. Magee LA, Pels A, Helewa M, Rey E, von Dadelszen P. Diagnosis, evaluation, and management of the hypertensive disorders of pregnancy: executive summary. *Journal of obstetrics and gynaecology Canada : JOGC = Journal d'obstetrique et gynecologie du Canada : JOGC*. 2014;36(7):575-6.

14. Xydopoulos G, Perry H, Sheehan E, Thilaganathan B, Fordham R, Khalil A. Home blood-pressure monitoring in a hypertensive pregnant population: cost-minimization study. *Ultrasound in Obstetrics & Gynecology*. 2019;53(4):496-502.

15. Hinton L, Tucker KL, Greenfield SM, Hodgkinson JA, Mackillop L, McCourt C, et al. Blood pressure self-monitoring in pregnancy (BuMP) feasibility study; a qualitative analysis of women's experiences of self-monitoring. *BMC pregnancy and childbirth*. 2017;17(1):427.

16. Carter W, Bick D, Mackintosh N, Sandall J. A narrative synthesis of factors that affect women speaking up about early warning signs and symptoms of pre-eclampsia and responses of healthcare staff. *BMC pregnancy and childbirth*. 2017;17(1):63.

17. Dougall D. FM, Tucker K., Yu L-M., Hinton L., Rivero-Arias O., Abel L., Allen J., Band R., Chisholm A., Crawford C., Green M., Greenfield S, Hodgkinson J., Leeson P., McCourt C., Mackillop L., Nickless A., Sandall J., Santos M., Tarassenko L., Velardo C., Wilson H., Yardley L., Chappell LC., McManus RJ. . Blood pressure monitoring in high risk pregnancy to improve the detection and monitoring of hypertension (The BUMP 1&2 trials): protocol for two linked randomised controlled trials. *BMJ Open*. 2019.
18. Lovell H, Wilson HM, Whybrow R, Harding K, Chappell LC. Caring for women with chronic hypertension. *Br J Midwifery*. 2019;27(6):343-51.
19. Hamilton A. Qualitative Methods in Rapid TurnAround Health Services Research VA HSR&D Cyberseminar Spotlight on Women's Health 2013 26.06.2017. Available from: [https://www.hsr.d.research.va.gov/for\\_researchers/cyber\\_seminars/archives/780-notes.pdf](https://www.hsr.d.research.va.gov/for_researchers/cyber_seminars/archives/780-notes.pdf).
20. Gale NK, Heath G, Cameron E, Rashid S, Redwood S. Using the framework method for the analysis of qualitative data in multi-disciplinary health research. *BMC medical research methodology*. 2013;13(1):117.
21. Ziebland S, McPherson, A. Making sense of qualitative data analysis: an introduction with illustrations from the DIPEX (personal experiences of health and illness. *Medical Education*. 2006;40(5):405-14.
22. Ltd. QIP. NVIVO qualitative data analysis software. 2018.
23. Mason M, editor Sample size and saturation in PhD studies using qualitative interviews. *Forum Qualitative Sozialforschung/Forum: Qualitative Social Research*; 2010.
24. Dougall G, Franssen M, Tucker KL, Yu L-M, Hinton L, Rivero-Arias O, et al. Blood pressure monitoring in high-risk pregnancy to improve the detection and monitoring of hypertension (the BUMP 1 and 2 trials): protocol for two linked randomised controlled trials. *BMJ open*. 2020;10(1).
25. Band R, Hinton L, Tucker KL, et al. Intervention planning and modification of the BUMP intervention: a digital intervention for the early detection of raised blood pressure in pregnancy. *Pilot Feasibility Stud*. 2019;5(153).
26. National Institute for Health and Care Excellence N. Hypertension in pregnancy: diagnosis and management NICE guideline [NG133]. 2019.
27. Jones MI, Greenfield SM, Bray EP, Baral-Grant S, Hobbs FR, Holder R, et al. Patients' experiences of self-monitoring blood pressure and self-titration of medication: the TASMINH2 trial qualitative study. *British Journal of General Practice*. 2012;62(595):e135-e42.
28. Hodgkinson JA, Tucker KL, Crawford C, Greenfield SM, Heneghan C, Hinton L, et al. Is self monitoring of blood pressure in pregnancy safe and effective? *BMJ*. 2014;349:g6616.
29. Johanson R, Newburn M, Macfarlane A. Has the medicalisation of childbirth gone too far? *BMJ*. 2002;324(7342):892-5.
30. McCourt C, Stevens T. Continuity of carer—what does it mean and does it matter to midwives and birthing women? *Canadian Journal of Midwifery Research and Practice*. 2006;4(3):10-20.
31. Bunton R, Petersen, A. Foucault, Health and Medicine. London: Routledge; 2002.
32. Wiltshire G, Fullagar S, Stevenson C. Exploring parkrun as a social context for collective health practices: running with and against the moral imperatives of health responsabilisation. *Sociology of health & illness*. 2018;40(1):3-17.
33. Newman CE, Persson A, Miller A, Brown RJ. "Just take your medicine and everything will be fine": Responsibilisation narratives in accounts of transitioning young people with HIV into adult care services in Australia. *AIDS care*. 2016;28(1):131-6.
34. Liebenberg L, Ungar M, Ikeda J. Neo-liberalism and responsabilisation in the discourse of social service workers. *The British Journal of Social Work*. 2013;45(3):1006-21.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

35. Garrett PM. How to be modern: New Labour’s neoliberal modernity and the Change for Children programme. *British Journal of Social Work*. 2006;38(2):270-89.

36. Rogers A, Kirk S, Gately C, May CR, Finch T. Established users and the making of telecare work in long term condition management: implications for health policy. *Social science & medicine*. 2011;72(7):1077-84.

37. Cohn S. From health behaviours to health practices: an introduction. *Sociology of health & illness*. 2014;36(2):157-62.

38. May C, Finch T, Mair F, Ballini L, Dowrick C, Eccles M, et al. Understanding the implementation of complex interventions in health care: the normalization process model. *BMC health services research*. 2007;7(1):148.

39. Ferrinho P, Sidat M, Goma F, Dussault G. Task-shifting: experiences and opinions of health workers in Mozambique and Zambia. *Human Resources for Health*. 2012;10(1):34.

40. Dawson AJ, Buchan J, Duffield C, Homer CS, Wijewardena K. Task shifting and sharing in maternal and reproductive health in low-income countries: a narrative synthesis of current evidence. *Health policy and planning*. 2013;29(3):396-408.

41. Bohmer RM, Imison C. Lessons from England’s health care workforce redesign: no quick fixes. *Health Affairs*. 2013;32(11):2025-31.

42. Mackintosh NJ. Interventions to increase patient and family involvement in escalation of care for acute life-threatening illness in community health and hospital settings. 2017.

43. Wolfe I, Cass H, Thompson MJ, Craft A, Peile E, Wieggersma PA, et al. Improving child health services in the UK: insights from Europe and their implications for the NHS reforms. *Bmj*. 2011;342:d1277.

44. Draper ES, Kurinczuk JJ, Kenyon S. MBRRACE-UK Perinatal confidential enquiry: term, singleton, normally formed, antepartum stillbirth: Infant Mortality and Morbidity Studies; 2015.

45. Warland J, O’Brien LM, Heazell AE, Mitchell EA. An international internet survey of the experiences of 1,714 mothers with a late stillbirth: the STARS cohort study. *BMC pregnancy and childbirth*. 2015;15(1):172.

46. Schwappach DL. Engaging patients as vigilant partners in safety: a systematic review. *Medical Care Research and Review*. 2010;67(2):119-48.

47. Intercollegiate Stroke Working Party. Public Report for England, Wales and Northern Ireland. National Sentinel Stroke Clinical Audit 2010. 2010.

48. Rance S, McCourt C, Rayment J, Mackintosh N, Carter W, Watson K, et al. Women's safety alerts in maternity care: is speaking up enough? *NBMJ Quality & Safety*. 2013;22(4):348-55.



TABLE 2 QUOTATIONS

THEME	QUOTATIONS
i) Different blood pressure changes in pregnancy	<p><b><i>BP Fluctuations</i></b></p> <p><i>"even as midwives we get confused with what should be normal because we have a protocol that they have a booking [.....measurement] so we've got a start and if it's 15 either way for the diastolic and we have to sort of think well yes that's creeping up a little bit if these girl's sent them to me I would look at it and say well that's not particularly high blood pressure because I'm dealing with PET all the time, you know, high blood pressure to me is 150/90 and above. So, isn't it, it's quite difficult and I think for a midwife let alone a woman to sit at home and think is that high am I going to be bothered"</i> <b>West Middlesex midwife FG3</b></p> <p><i>"Another thing outside of pregnancy and not, most of what you'll doing with hypertension monitoring is trying to prevent long term complications on a short term and therefore, actually what you, slightly over 24 hours probably is what really matters whereas in pregnancy [...] I'm actually only interested in whether or not they've got [...] Well actually I'm really interested in the next 24 hours."</i> <b>Guys and St Thomas's obstetrician FG1</b></p> <p><b><i>Wider symptoms</i></b></p> <p><i>"Yes but I think you raise a really important point about when you start to kind of focus on individual symptoms of a condition it's, there's a risk that you just focus on the blood pressure which means if you're feeling dizzy or your ankles are puffy or you're not really feeling yourself and got a blinding headache you don't focus on any those things because the blood pressure says it's okay."</i> <b>Royal London, Community midwife</b></p> <p><i>"I think that they're over checking their blood pressure and occasionally I think it is helpful that they see a midwife and do things like check the protein in the urine and things like that"</i></p>

	<p><i>which, you know, occasionally perhaps not be able to manage quite so well yourself</i> <b>John Radcliffe, Obstetrician</b></p> <p><i>"So you kind of think oh are we only doing half of a job if they're just monitoring their blood pressure"</i> <b>Horton, Community Midwife</b></p>
ii) Reliability and accuracy	<p><i>"As a health professional I know blood pressure's affected by many things, whether you are taking it standing whether you are taking it lying whether it's in the morning whether you, or you've just come back from Tesco with plastic bags and stuff like, you're out of your breath or you've just fought with your partner or something like that, so many things influences blood pressure. So they will need to be given the empowerment they will need to take their blood pressure so that we, it's as accurate as, so that we get as close to the normal if that can be achieved, something like that."</i><b>West Middlesex midwife FG2</b></p> <p><i>"[R4] . Because there used to be a particular, yes because women's bodies are totally different in pregnancy to somebody having a blood pressure normally and so, you know, for me I've never looked into it but whether it's a 100% accurate I don't know but I certainly don't like those box things because they have to be quiet, silent and not moving and nobody talking to them.</i></p> <p><i>[R1]: That's why the machine is important.</i></p> <p><i>[R4]: To get a really accurate result."</i><b>Whipps Cross, FG1</b></p> <p><i>"I wouldn't ignore it if it was very elevated but I would certainly repeat it, I wouldn't go with what they said... Because I'd be worried about the quality of their machine"</i> <b>John Radcliffe, Obstetrician</b></p> <p><i>"we advise them not to because [um] they're not calibrated, you know, they're their own there's no way they're calibrated and also quite a lot of [um] blood pressure [um] equipment, we've read research that says it's not as accurate as manual or a professional taking the blood pressure"</i> <b>Royal Berkshire, FG1</b></p>

	<p><i>"when they come along and they bring readings [um] and they, if there's any conflict and sometimes they don't want to be labelled as hypertensive because it leads to certain choices later on in pregnancy that may be removed from them i.e. where they deliver, how they deliver [um] I then, you know, I will talk to them and say you, whatever you're using at home may or may not be validated for pregnancy but what we use here is validated for pregnancy and therefore I would defer to those readings"</i> <b>Royal Wolverhampton, Consultant Obstetrician</b></p> <p><i>"I tend to be a little bit sceptical because you don't know what they're using, you don't know what machine they're using, you don't know whether they're trained to use it, you don't know how old that machine is whether it's again cuff size, you know, if they've borrowed it off somebody you don't know whether it's been PAT tested, you know, and if they are concerned about their reading being particularly high or particularly low I'd rather just get them in to see the relevant person at the time rather than rely on that, you know, I do worry about them using their own equipment because you just don't know how accurate it is"</i> <b>Royal Wolverhampton, FG2</b></p> <p><i>"I'd want to know whether or not the machine was up to scratch xxx good if she could correlate it with ours, bring it in we'll just make sure your machine is similar to ours, there's things like that."</i> <b>Royal Wolverhampton FG3</b></p>
<p><b>iii) Impact on women</b></p>	<p><i>"Those people with raised blood pressure will share their experience of like three or four weeks admission into hospital I think they will be delighted to do it to save coming in, they always, you know, get very upset having to stay on for three weeks and when they have child at home."</i> <b>West Middlesex FG3</b></p> <p><i>"So for the women who have had severe pre-eclampsia in the past who are absolutely panic stricken about it being missed in the future [.....] I don't know scientifically if it matters but for them it gives them huge reassurance that they've got, it means that rather than them having to see the clinician every week or</i></p>

	<p><i>twice a week, they're doing something, it's rather like feeding a baby every other day."</i> <b>Guys and St Thomas's FG1</b></p> <p><i>"There's a lot of women that have mildly raised blood pressure that are fine, that we don't really do anything with. [Um] and it's a lot of monitoring for them"</i> <b>John Radcliffe, Obstetrician</b></p> <p><i>"some women might not want to monitor themselves because they wouldn't feel, it will be too much responsibility in case they miss something"</i> <b>John Radcliffe FG3</b></p> <p><i>"I think it's a bit of mix isn't it, I think some women would really like it saves them time especially if they're busy they have other children."</i> <b>John Radcliffe FG3</b></p> <p><i>"I think a lot of women would also appreciate it because they often don't want to come into hospital and they would, I think if they were self-monitoring at home if they're taught in the right way and they know when somethings high or they know when to escalate or to ring and I think it will work really well and stop a lot of people coming in unnecessarily. I think it would be really good"</i> <b>John Radcliffe, Hospital Midwife</b></p>
<p><b>iv) Anticipated impact on the antenatal care</b></p>	<p><i>"we are so busy, we are too busy in hospitals and the more we can do out of hospital safely the better"</i> <b>John Radcliffe, Obstetrician</b></p> <p><i>"there are benefits all-round aren't there, if you're, if we're [um] happy that this woman is monitoring her blood pressure sensibly [um] it's got benefits for her in that it means she doesn't have to come so often, it's got benefits for us in that it reduces our, our workload, surely it's got, it has a, there's an economic benefit there for the NHS [um] as a whole and it, I suppose that's one of the things isn't it that public health it's about educating people isn't it about, so that, you know, I think there are lots of benefits it's [um] it's just getting it in place. And being confident about the thing that you've got in place"</i> <b>Whipps Cross, Day Assessment Unit midwife</b></p> <p><i>"I think it would have the potential to increase the appointments rather than decrease appointments as a screening tool for</i></p>

	<p><i>hypertension because of the worried well who've seen systolic increase in ten as an example but I don't think it will decrease the routine ante-natal screening because that has many other roles including measurement of fundal height or foetal wellbeing."</i> <b>Royal London, medical trainee</b></p> <p><i>"I suppose potentially it will increase workload because people are going to recognise that they have a slightly elevated blood pressure sooner but I don't think that's necessarily a bad thing"</i> <b>John Radcliffe, Obstetrician</b></p> <p><i>"So it's a massive information imparting exercise probably more importantly than the, the systematic screening for obstetric complications that old fashioned people think of as ante-natal care. You know I, you know, again when I trained we'd see clinics of 40 people, 50 people in a morning where literally all you do would be check their blood pressure, check their urine, put your hand on their tummy, check their heart, you know, see if the patients still breathing, next, you know. But it isn't like that now there's a lot more education, seeing much more as a health education exercise as much as a screening for abnormality of pregnancy exercise".</i> <b>Royal Berkshire, Consultant Obstetrician</b></p>
<p><b>v) Caution, uncertainty and evidence</b></p>	<p><b>Caution</b></p> <p><i>"I wouldn't take the, take everything as a 100% as gospel because, you know, you can't really it's like triaging over the phone they can tell you so much but until you've seen that woman and made an assessment you can't do a full assessment from that. But [um] but yes I mean it would certainly I, I like compare to a jigsaw puzzle, you know, putting all the pieces together that would definitely be one of the pieces I would definitely include once I knew that all the bits were safe that they were using and everything, yes."</i> <b>Royal Wolverhampton FG2</b></p> <p><i>"if I look at it and I can see that yes at 2:00 'o' clock and half two she had raised blood pressure but [...] guidelines about what I do about that [um] all I can do is do my own blood pressure readings and get them reviewed by the doctor. Unless there was</i></p>

	<p><i>like a real, you know, like a guideline of what we actually do about the trends, there's not much I myself could actually do from it.</i> "Royal Wolverhampton FG4</p> <p><b>Uncertainty</b></p> <p><i>"I think that would be very useful information to have rather than as you say the snap shot that you get in clinic. Although on the opposite side if they were having consistently low readings at home and when they come and visit us in triage it's always 160/110 I would question whether the data was true even if I knew that the machine was valid. Acknowledging the role of white coat hypertension but I would still be very worried about relying purely on the home readings and discarding the readings I have in the hospital."</i> <b>Royal London, medical trainees.</b></p> <p><i>"You can also have high blood pressure at home and lower blood pressure readings in hospital and we have no idea, I don't think [um] what we should do when actually readings are different in different places and there is a tendency, there will be some people who will have a tendency to treat anybody whatever the excuse, much more common is a tendency to say oh it's probably more, we've had 150, 140, 120 I'm going to believe the normal whether its home reading or hospital readings. And so just having more data may just be making more noise rather than giving us, telling us what to do. But without the data we don't even know if that's true."</i> <b>Guys and St Thomas's FG1</b></p> <p><i>"supposed to be evidence based and when you actually look at them about 90% of the recommendations they make are expert opinion"</i> <b>Royal Berkshire, Consultant Obstetrician</b></p>
vi) Concerns over action/inaction	<p><i>"Yes exactly and they may not realise the seriousness of it, so that would be my concern if they're doing it at home. The chronic hypertension women have a lot of contact with, in hypertension clinic they understand blood pressure they've been dealing with it, they're on medication they understand something about blood pressure, they might not fully understand how serious it can be</i></p>

*with regards to pre-eclampsia but they certainly understand that they need to act on it and if they don't, obviously it's explained to them numerous times. My concern is if you did that to a general population will that message be translated."* **Guys and St Thomas's FG2**

*"I do have a worry about women taking, you know, something like pre-eclampsia is a multi-system disorder, blood pressure is one component and we see every week, every month women who come in with normal blood pressure but everything else going wrong and it would make me nervous about the idea of women taking control of their care so much so that they felt reassured by one reading and ignored their signs and symptoms and I think that, that will be, that's a major, that's a clinical worry that I have for using."* **John Radcliffe, Consultant Obstetrician**

*"I think as long as they [women] understand the limits of what's normal and what's not and they know the right people to contact if not, then I don't see there's anything wrong with it"* **Royal Berkshire FG2**

*"I just know that for some women they, they've got so much going on that actually their health is, is quite easily overlooked and [um] is last on the agenda if that makes sense and if they could miss a blood pressure reading here and there [um] then they may well do that and that would be my only [um] and which I feel awful saying really because I am so like passionate about actually women being responsible for their own care and taking ownership of it and us giving it back to them [um]. [.....] I just think some women will be like oh yes, everything's fine they're all okay and maybe overlook things. I know you've got the opposite extent where some people would potentially act on it like say there's a higher reading because they think it will be acted on and that sort of thing as well but I think you could almost get women who will either pretend they've done it or pretend that it's an okay reading when it's not [um] which is awful to say that we don't trust women but"* **Royal Wolverhampton FG2**



**SELF MONITORING OF BLOOD PRESSURE IN PREGNANCY**

**(WS1.2 FOCUS GROUPS WITH STAFF)**

**INTRODUCTION**

*SCRIPT: e.g. “we are interested in finding out if asking women to monitor their own blood pressure during pregnancy, in addition to routine antenatal monitoring, can improve the early detection of hypertension and improve their care. We would like to hear your views on this, and would like to find out how we might best introduce women’s home readings into current care pathways”*

**QUESTIONS**

**(a) Attitudes to self-monitoring**

In general, what are your attitudes to women taking their own blood pressure readings at home?

What do you think about the accuracy of home readings? Would you be prepared to take action on the basis of home readings (either normal or high)?

How do you think taking BP readings at home would make women feel?

Do you have a view on how many days per week women should monitor their BP? (3 or 4)

How would such SMBP readings be handled within standard antenatal care appointments? Would you record them in the patient record?

What would you do next if a woman presented with raised BP at home?

If SMBP readings were not congruent with those in the clinic, what would you do (e.g. high at home, normal in clinic or vice versa)?

What approach to BP monitoring do you trust the most and least – clinic, home or ABPM?

What do you think the impact of more women self-monitoring might be on usual maternity care? Do you think there would be any impact on workloads?

**(b) Outcomes**

What data would be most influential in changing your practice to include self-monitoring? [earlier diagnosis, better BP control, reduced complications]

**(c) Operationalising self-monitoring**

What would be the best way to integrate SMBP readings into antenatal care in your hospitals?



Are there any barriers to introducing home readings from women into these (antenatal care) pathways?

What would help integrate these readings into antenatal care?

Have you any thoughts as to how we can support women who are self-monitoring to continue with it? What sort of training do you think women will need?

How can we make recruitment work well for staff and women? (we plan to approach women at 12 weeks with info, recruitment into study at 20 weeks – do you think this is sensible [note early drop out in pilot with miscarriage etc])

How do you think we should identify women who are taking part in the notes etc? What information would you like to see in the notes about the study? How would you feel about a study “passport” that alerts you to women taking part?

Who should have access to the BP data women send in via the web-based management system?

What would be the best way to deliver training for you/ staff? (face to face, web based etc) Do you feel you have any particular training needs/ What needs to be clearer?

**PI Interviews**

**(a) Current practice**

- Check the terminology used for clinics in each of the sites (e.g. maternity assessment unit/MAU etc)
- What is your involvement in BP monitoring of pregnant women?

How is blood pressure monitoring of pregnant women (and identification of hypertension) currently handled in your hospital?

- Who generally does the measurements?
- How often is BP measured, typically? What would be a minimum number of times BP is measured? What would be the maximum?
- What monitors are used? Are they validated? How often are they calibrated?
- Is ABPM used?
- Who is the BP result shared with in your hospital/ Who has access to the record?

If raised BP is detected in clinic, what happens next to women?  
[for example, primagravida at 30/40 with BP 149/95 in clinic, no protein]

- What additional tests or monitoring would they undergo?
- At what BP level would treatment be offered?

Does the hospital ever encourage SMBP/ consider home BP data?

**(b) Operationalising self-monitoring**

What would be the best way to integrate SMBP readings into antenatal care in your hospitals?

Are there any barriers to introducing home readings from women into these (antenatal care) pathways?

What would help integrate these readings into antenatal care?

Have you any thoughts as to how we can support women who are self-monitoring to continue with it? What sort of training do you think women will need?

How can we make recruitment work well for staff and women?

How do you think we should identify women who are taking part in the notes etc?

What information would you like to see in the notes about the study?

How would you feel about a study “passport” that alerts you to women taking part?

Who should have access to the BP data women send in via the web-based management system?

Who should we talk to, to make accessing the web based system from the wards/ clinic as easy as possible?

For peer review only